

WellBeing International

WBI Studies Repository

2007

An HSUS Report: Human Health Implications of U.S. Live Bird Markets in the Spread of Avian Influenza

The Humane Society of the United States

Follow this and additional works at: https://www.wellbeingintlstudiesrepository.org/hsus_reps_environment_and_human_health



Part of the [Animal Studies Commons](#), [Other Animal Sciences Commons](#), and the [Veterinary Infectious Diseases Commons](#)

Recommended Citation

The Humane Society of the United States, "An HSUS Report: Human Health Implications of U.S. Live Bird Markets in the Spread of Avian Influenza" (2007). *Impact of Animal Agriculture*. 9.

https://www.wellbeingintlstudiesrepository.org/hsus_reps_environment_and_human_health/9

This material is brought to you for free and open access by WellBeing International. It has been accepted for inclusion by an authorized administrator of the WBI Studies Repository. For more information, please contact wbisr-info@wellbeingintl.org.



An HSUS Report: Human Health Implications of U.S. Live Bird Markets in the Spread of Avian Influenza

Abstract

After millions of years of existing as a harmless infection of aquatic birds, some strains of avian flu have developed the ability to mutate into highly pathogenic forms that may not only be deadly for birds, but potentially more dangerous for humans as well. Experts from the World Health Organization, the World Organization for Animal Health, and the Food and Agriculture Organization of the United Nations have joined leading scientists in implicating industrialized poultry production for providing fertile ground for the transformation of benign bird flu into highly pathogenic strains.

Live bird markets can be the conduit by which waterfowl viruses enter into industrial poultry populations and have been identified as playing a critical role in the emergence of H5N1, a deadly strain of avian influenza threatening to trigger a human flu pandemic. In response, Asian countries are shutting them down. In the United States, live bird markets have been tied to domestic outbreaks of highly pathogenic avian influenza and have been described by U.S. Department of Agriculture poultry researchers as the “missing link in the epidemiology of avian influenza,”¹ yet hundreds of these retail storefront slaughter markets still operate across the country, processing tens of millions of birds annually.

Background on Avian Influenza

The primordial source of all influenza viruses—avian and mammalian—is aquatic birds.² In nature, the influenza virus has existed for millions of years as a harmless, intestinal, waterborne infection of waterfowl, particularly ducks,³ who are infected as ducklings. Studies of ducks on Canadian lakes show that up to 30 percent of juvenile birds are actively shedding the virus.⁴ The birds excrete such massive titers of virus⁵ that researchers have been able to culture it straight out of a spoonful of lake water.⁶ Under the right conditions, the virus is estimated to be able to persist for years in cold water.⁷ With such high concentrations of virus and its highly efficient transmission⁸ and environmental stability, scientists estimate that virtually all of the millions of ducks in the world become infected sometime during their lives.⁹

Avian influenza multiplies in the ducks’ intestinal lining and is then excreted into the water of ponds, lakes, and other reservoirs. Ducks touching down to drink ingest the virus, and the cycle continues as it has for perhaps 100 million years before the first person became infected with the flu.¹⁰ The ducks are not infected for long;¹¹ most only shed virus for a few days. However, the fecal-oral route of infection for aquatic birds is so efficient that this narrow period of contagion is thought to be enough to keep the virus spreading throughout the millennia.¹² Millions of years of evolution have so tailored the parasite to its host, a so-called “optimally adapted system,” that the virus seems completely innocuous to the ducks.¹³ The virus exists in an “evolutionary stasis” in waterfowl,¹⁴ remaining unchanged despite its rapid mutation rate.¹⁵

Influenza viruses do not tend to retain any new mutations in ducks because they found their perfect niche. Sharks, for example, have remained basically unchanged for 100 million years, even while other species evolved around them, presumably because sharks had already evolved to be such perfected predators.¹⁶ Similarly,

evolution has not seemed able to significantly improve influenza. While the virus continues to mutate to achieve greater and faster spread, the virus-duck relationship seems so cooperative that any deviant viral progeny are less successful and die out. The virus has achieved peak efficiency and is found ubiquitously throughout aquatic bird species around the world.¹⁷

As the influenza virus in its natural state is so finely attuned to its aquatic host, it not only doesn't harm the carrier duck, but it also seems unable to cause serious disease in humans. Before the emergence of H5N1, the deadly mutant strain of avian influenza spreading out of Asia, there had been only two reports of human infection from wild bird viruses in the medical literature: one involving a woman who kept ducks and got a piece of straw caught in her eye while cleaning out her duck house,¹⁸ and the second of a laboratory field worker whose eye was sneezed into by a bird flu-infected seal.¹⁹ Despite direct inoculations of virus, the result was limited to a mild, self-contained (unspreading) case of conjunctivitis, commonly known as pinkeye.²⁰

Scientists have been relatively unsuccessful in attempts to directly infect human volunteers with waterfowl viruses. Test subjects snorted massive doses of virus—enough to infect up to a billion birds—yet, most of the time, did not become infected themselves. When the virus did have an effect, it typically produced nothing more than a transitory local reaction.²¹ The evolutionary gulf between duck gut and human lung is so broad that an intermediate host is thought to be needed to act as a stepping stone for the virus.²² “And poultry,” said a spokesperson for the Department of Homeland Security’s National Center for Foreign Animal and Zoonotic Disease Defense, “are likely to be that host.”²³ Bird flu viruses only “heat up,” in the words of Dutch virologist Albert Osterhaus, “when they pass from wild birds to poultry.”²⁴

Mutation

All bird flu viruses begin as mild, low-grade, low-pathogenicity avian influenza (LPAI) viruses. H5 and H7 virus strains, however, have the potential to mutate into high-grade “fowl plague” viruses, now known as HPAI—highly pathogenic avian influenza.

In its 2005 assessment of the pandemic threat, the World Health Organization explained the emergence of HPAI viruses: “Highly pathogenic viruses have no natural reservoir. Instead, they emerge by mutation when a virus, carried in its mild form by a wild bird, is introduced to poultry. Once in poultry, the previously stable virus begins to evolve rapidly, and can mutate, over an unpredictable period of time, into a highly lethal version of the same initially mild strain.”²⁵

Live bird markets have been identified as enabling the introduction of low-grade, waterfowl avian influenza viruses to commercially reared poultry. Consider the following scenario: An infected duck is transported to a live poultry market where a variety of bird species are typically confined in cages stacked in tiers. The duck’s droppings fall outside the wire enclosure onto other birds, customers and workers, and the surrounding environment. Direct human contact with the infected feces may not result in the unmutated duck virus taking hold. However, terrestrial birds, such as quail and chickens, two species commonly found at live bird markets, are not natural hosts for influenza,²⁶ yet may be recognizable enough by the virus to become infected. Despite this, the influenza virus, finely attuned to ducks and other waterfowl and having remained in total evolutionary stasis in aquatic birds,²⁷ must mutate to survive and effectively spread in, for example, the gut of a chicken. The influenza can no longer remain a strictly waterborne virus. Although it can still spread through feces when chickens peck at each other’s droppings, the virus must, for example, resist dehydration better in the open air²⁸ and presumably adapt to the novel body temperature and pH of its new environment.²⁹

In ducks, the virus relies on a healthy host for its spread and seems to have evolved a mechanism that allows the virus to replicate only in the intestine, so as not to infect other tissues and potentially harm the carrier. However, in such foreign environments as land-based poultry, influenza—highly effective at mutating³⁰—quickly begins to try to adapt to the new host³¹ in order to survive.^{32,33} Given adequate time and numbers of hosts within

which to mutate, some bird flu viruses are able to learn how to invade other organs, such as the lungs, through which to travel.

Evolved Harmlessness and Transitioning Virulence

Before the virus can become infectious, a protein spike in its viral coat called hemagglutinin (the “H” in H5N1) must first be cleaved in half to become activated. This cleavage is done by specific host enzymes found only in certain tissues, including the intestinal tract. The limited bodily distribution of the specific cleavage enzyme restricts the virus to safe areas such as the duck’s gut, preventing it from replicating in the brain or other vital organs. This restriction, evolved harmlessness, has allowed influenza to most effectively pass on and spread its genes.³⁴ Yet, once the virus successfully infects an unfamiliar species, this may no longer be the case. Researchers have shown that H5N1 seemed to enter chicken populations as an intestinal virus but left as more of a respiratory virus.³⁵

Once a foreign host is infected, the virus must not only find new ways to spread; it must also overcome a hostile immune response. Under these circumstances, viral progeny that eliminate the fail-safe mechanism that restricts it to the gut may outspread others and be naturally selected. The hemagglutinin spike of H5 and H7 viruses can mutate over time to be activated by enzymes in any organ in the body, allowing the virus to essentially liquefy the bird from the inside—the “flubola” phenomenon of highly pathogenic avian influenza.³⁶ While influenza remains harmless to its waterfowl hosts in its natural state, the virus can be forced into what evolutionary biologists call an “acute life strategy” in which its only choice may be to rapidly overwhelm the host to gain entry and survive in new environments.³⁷ Viral mutants with diminished lethality may have an overall advantage over strains that kill the host too quickly, thereby limiting the opportunities for infectious spread, since the host may stay alive longer and have more occasion to pass on the virus. Natural selection mediates this evolutionary process, choosing over time the virus with the perfect balance of lethality and contagion.³⁸ However, transitions are not always toward lesser virulence.³⁹

Infecting an unfamiliar host such as a chicken, the influenza virus may become as virulent as possible to overpower the bird’s defenses.⁴⁰ In a natural setting, if the virus kills the chicken too quickly, it may not be able to infect many others. Yet, in unnatural settings such as those common in intensive poultry production where tens of thousands of genetically similar birds are intensively confined in a single enclosure, there may be fewer limits to how virulent certain influenza viruses can become and still survive.

Mutation Facilitation by Industrial Poultry Production

Scientists have demonstrated the transformation of viruses from harmless to deadly in a laboratory setting. A collaboration of U.S. and Japanese researchers started with H5N3, a harmless virus isolated from a whistling swan, unable to infect a healthy, adult chicken. The researchers proceeded to do serial passages through baby chickens, squirting a million infectious doses into the lungs of day-old chicks. The virus would presumably start to adapt to the chicks’ respiratory tracts, with the viral mutant that most effectively and efficiently overcame the hatchlings’ defenses selected to predominate. After three days, the scientists killed the chicks, ground up their lungs, and introduced this infectious material down the throats of other chicks. They again allowed a few days for the virus to adapt further before repeating the cycle two dozen times.

When the last set of chicks was killed, the researchers ground up their brains instead of their lungs and infected five additional rounds of healthy chicks with the brain pulp. With every passage, the virus grew more adept at overwhelming the fledging birds’ immune systems to best survive and thrive in its new environment. After two dozen cycles through lungs and five cycles through brain, the final infected brain sample was squirted into the nostrils of healthy, adult chickens. Although the original H5N3 swan virus was unable to infect the chickens, by the 18th lung passage, the strain was able to kill half of the chickens exposed. After the final five brain passages, the virus was capable of rapidly killing every one. The researchers concluded: “These findings demonstrate that

the avirulent [harmless] avian influenza viruses can become pathogenic during repeated passaging in chickens.”⁴¹

Industrial poultry production settings allow for the potential of tens of thousands of such serial passages in a single building. The World Organization for Animal Health (OIE) and the Food and Agriculture Organization of the United Nations (FAO) agree that it has been “prove[n]”⁴² that once LPAI viruses gain access to poultry facilities, they can “progressively gain pathogenicity in domestic birds through a series of infection cycles until they become HPAI.”⁴³ However, deadly bird flu viruses do not tend to arise in merely any poultry operation. According to U.S. Department of Agriculture researchers, it is the “high density confinement rearing methods” that give bird flu “a unique chance to adapt to the new species.”⁴⁴ That is, today’s intensive animal production practices may remove the natural obstacles to transmission that prevent the virus from becoming too dangerous.

Director of the USDA’s chief poultry research laboratory and the agency’s leading bird flu expert, David Swayne has authored more than 100 scientific publications on avian influenza.⁴⁵ According to Swayne, there has never been a recorded emergence of an HPAI virus in any backyard flock or free-range poultry operation. If a wild duck were to excrete a dropping laden with a relatively innocuous virus into a free-range flock, the chickens may be exposed to the virus, but coming straight from waterfowl, the virus is so finely tuned to duck physiology that it may not gain a foothold before being killed by a healthy chicken’s immune system. This is why in the laboratory, researchers *injected* infected lung tissue from one bird to another to facilitate transmission. “The conditions under which we generated highly virulent viruses from an avirulent strain are generally not duplicated in nature,” the research team acknowledged. “However, viruses with low pathogenicity can cause viremia in physically compromised chickens.”⁴⁶ Viremia means successful invasion of the bloodstream by the virus, an incursion they deem more likely to occur in compromised hosts.

If an outdoor flock does become infected, the virus must continue to spread to remain in existence. Influenza virus, a waterborne virus, is rapidly killed by sunlight and tends to dehydrate to death in the breeze. Its ability to spread efficiently from one chicken to the next while outdoors in the open air is relatively limited. In a sparsely populated free-range setting, there may simply be too few susceptible hosts nearby to pass between in order to build up enough adaptive mutations to evolve into a highly pathogenic form. According to bird flu expert Dennis Alexander of the U.K.’s Central Veterinary Laboratory, highly pathogenic influenza viruses have “never known to arise in an outdoor [chicken] flock.”⁴⁷

In contrast, the conditions and practices of industrial chicken production are considered to be optimal for dangerous viral mutations.⁴⁸ Industry trade journal *World Poultry* listed some factors that make intensive poultry facilities such “ideal”⁴⁹ “breeding grounds for disease”⁵⁰: “poor ventilation, high stocking density, poor litter conditions, poor hygiene, high ammonia level, concurrent diseases and secondary infections.”⁵¹ Indeed, an avian virology textbook states: “Viral infections can move fastest through groups of birds maintained in closed, crowded, unsanitary conditions.”⁵² Frederick Murphy, dean emeritus of the School of Veterinary Medicine at the University of California-Davis, has noted how intensification in farm animal production practices “often allow[s] pathogens to enter the food chain at its source and to flourish, largely because of stress-related factors.”⁵³ The physiological stress created by crowded confinement can have a “profound” impact on immunity,⁵⁴ predisposing animals to infection.⁵⁵ Diminished immune function reduces protective responses to vaccinations. “As vaccinal immunity is compromised by factors such as... immunosuppressive stress,” writes a leading⁵⁶ USDA expert on chicken vaccines, “mutant clones have an increased opportunity to selectively multiply and to be seeded in the environment.”⁵⁷ Studies exposing birds to stressful housing conditions provide “solid evidence in support of the concept that stress impairs adaptive immunity in chicken.”⁵⁸

Typical broiler chicken production practices entail intensively confining tens of thousands of genetically undiverse⁵⁹ birds in massive, unhygienic warehouses, standing and lying in their own excrement. The high stocking densities of commercial poultry production result in an extraordinary amount of waste, and as influenza can survive in wet manure for weeks, these unhygienic conditions pose significant risk. A 20,000-bird broiler

chicken flock produces more than a ton of droppings every day.⁶⁰ And, according to the USDA, a single gram of manure (approximately the weight of a paper clip) from an infected chicken can contain “enough virus to infect 1 million birds.”⁶¹

Moist fecal dust and ammonia in the air irritate the birds’ respiratory passages, further increasing susceptibility in chickens already compromised by the stress of confinement. Feces decomposition generates several irritating chemicals, including hydrogen sulfide, methane, and ammonia,⁶² which “in a poultry house is nauseating to the caretaker, irritates the eyes, and affects the chickens,” states one poultry science textbook.⁶³ Given the extreme stocking density of intensive production facilities, the litter can get so saturated with excrement that birds may develop sores or “ammonia burns” on their skin, known as breast blisters, hock burns, and footpad dermatitis, all of which have become significantly more common and serious over the last 30 years.⁶⁴

Studies have shown that high levels of ammonia also increase the severity of respiratory disorders, such as pneumonia,⁶⁵ in part by directly damaging the respiratory tract, predisposing birds to infection.⁶⁶ A large-scale study of millions of birds from nearly 100 commercial farms across multiple countries found that ammonia levels increased the excretion of the stress hormone corticosteroid, a potent immune depressant.⁶⁷

Since the birds are standing in their own excrement, the virus need not even develop true airborne transmission via nasal or respiratory secretions. Rather, the virus has an opportunity to be excreted in the feces and then inhaled or swallowed by the thousands of other birds confined in the shed, allowing the virus to circulate rapidly and repeatedly.

Industrial Poultry Production Implicated by Global Experts

The World Health Organization (WHO), the OIE, and the FAO, respectively the world’s leading medical, veterinary, and agricultural authorities, all implicate industrial poultry production as playing a role in the current avian influenza crisis. The WHO blames the increasing trend of emerging infectious diseases in part on the “industrialization of the animal production sector”⁶⁸ in general, and the emergence of H5N1 on “intensive poultry production” in particular.⁶⁹ The OIE blames in part the shorter production cycles and greater animal densities of modern poultry production, which result in a “greater number of susceptible animals reared per given unit of time.”⁷⁰ Said one senior FAO official, “[I]ntensive industrial farming of livestock is now an opportunity for emerging diseases.”⁷¹

Other experts around the world similarly lay blame at least in part on “so-called factory farming,”⁷² “intensive poultry production,”⁷³ “large industry poultry flocks,”⁷⁴ “intensive agricultural production systems,”⁷⁵ or “intensive confinement.”⁷⁶ “We are wasting valuable time pointing fingers at wild birds,” the FAO has stated, “when we should be focusing on dealing with the root causes of this epidemic spread which...[include] farming methods which crowd huge numbers of animals into small spaces.”⁷⁷

In October 2005, the United Nations issued a press release on bird flu specifically calling on governments to fight what they call “factory farming”: “Governments, local authorities and international agencies need to take a greatly increased role in combating the role of factory-farming, commerce in live poultry, and wildlife markets which provide ideal conditions for the virus to spread and mutate into a more dangerous form....”⁷⁸

Emeritus professor Kennedy Shortridge, the virologist who first characterized H5N1 in Asia, was awarded the highly prestigious Prince Mahidol Award in Public Health, considered the “Nobel Prize of Asia,”⁷⁹ for his pioneering work on H5N1.⁸⁰ From 1977 to 2002, he advised the World Health Organization on the ecology of influenza viruses.⁸¹ Shortridge described how modern poultry operations have created the greatest risk scenario in history. “The industrialization is the nub of the problem,” he said. “We have unnaturally brought to our doorstep pandemic-capable viruses. We have given them the opportunity to infect and destroy huge numbers of birds and...jump into the human race.”⁸² Johns Hopkins neurovirologist R.H. Yolken and Stanley Medical Research

Institute director E.F. Torrey agree: “In our efforts to streamline farming practices to produce more meat for more people, we have inadvertently created conditions by which a harmless parasite of wild ducks can be converted into a lethal killer of humans.”⁸³

The Role of Live Bird Markets

Analyzing the genome of H5N1, scientists now suspect that its original 1997 Hong Kong outbreak arose when an H5 goose virus combined with an N1 duck virus with quail acting as the mixing vessel. The virus then jumped from quail to chickens and then from chickens to humans.⁸⁴ Cross-contamination at live bird markets is the likely explanation of how these viruses jumped from species to species and, ultimately, farm to farm.

According to expert Professor Shortridge, live bird markets are an “avian influenza melting pot.”⁸⁵ Animals sold live guarantee freshness in the minds of some consumers, but the conditions of typical storefront slaughter facilities are unhygienic and severely compromise the animals’ well-being: animals cough and defecate over one another, spewing potential pathogens throughout the market;⁸⁶ chickens, ducks, geese, quail, and other birds are confined into restrictive cages stacked in tiers as much as five high, with distressed birds defecating on those below them;^{87,88} and feathers and feces,⁸⁹ as well as blood and intestines⁹⁰ soil the market. According to the WHO, the birds are often slaughtered on the spot, “normally with very little regard for hygiene.”⁹¹ “The activities of humans have affected the evolution of influenza,” reads a 2004 Cambridge University Press textbook on pathogen evolution, “but not to our advantage. Close confinement of various strains of fowl in live poultry markets provide conditions ripe for the formation of new reassortment viruses and their transmission to humans.”⁹² “If you have seen these markets, you know that the birds are under stressful conditions,” said a veterinarian with New York’s Department of Agriculture. “And birds under stress are much more prone to disease.”⁹³

Birds who remain unsold at the end of the day may go back to nearby farms, taking whatever new viruses they picked up with them.⁹⁴ Leading flu researchers Robert Webster and D. J. Hulse wrote, “Highly concentrated poultry and pig farming, in conjunction with traditional live animal or ‘wet’ markets, provide optimal conditions for increased mutation, reassortment and recombination of influenza viruses.”⁹⁵ Considering the Hong Kong outbreak and other outbreaks around the world traced to live poultry markets,^{96,97} USDA poultry researchers describe live bird markets as the “missing link in the epidemiology of avian influenza.”⁹⁸

Live Bird Markets in the United States

Live bird markets are not limited to Asia.⁹⁹ The USDA estimates more than 20 million birds of different species pass through 150 known storefront live bird markets just in northeastern U.S. metropolitan areas every year.¹⁰⁰ In addition, approximately 50 such markets exist in California¹⁰¹ and a half-dozen in each of the states of Minnesota,¹⁰² Texas,¹⁰³ and Florida.¹⁰⁴ Unlike many Asian countries, which have responded to the risks associated with live bird markets by shutting them down, and Hong Kong, which now segregates waterfowl from terrestrial species, U.S. live bird markets are still in operation, and separation of aquatic and land-based birds is not mandated.¹⁰⁵

Avian Influenza in the United States

Bird flu viruses have been detected every year in the United States since the mid-1960s.¹⁰⁶ In just the last five years, more than a dozen outbreaks of viruses with the potential to mutate into highly pathogenic forms have been discovered.¹⁰⁷

Prior to the emergence of the H5N1 strain of avian influenza in Hong Kong, the biggest outbreak of bird flu in history was in the United States in 1983—the nation’s most costly and extensive disease eradication. The outbreak of H5N2 avian influenza in Pennsylvania that led to the deaths of 17 million chickens,¹⁰⁸ spread down through Maryland and Virginia,¹⁰⁹ and cost taxpayers and consumers more than \$400 million,¹¹⁰ was considered

by some experts to have a link to live bird markets in the northeastern United States.¹¹¹ Before then, highly pathogenic bird flu had struck in the 1920s and later occurred in Texas in 2004. Both of these other high-grade bird flu incidents,^{112,113} as well as most of the latest U.S. discoveries of low-grade viruses, have also been linked to live poultry markets in the United States.¹¹⁴

Highly Pathogenic H5N2 and Live Bird Markets

Investigators speculate that the H5N2 virus responsible for the 1983 outbreak may have initiated in a flock of wild ducks who landed in a pond on a chicken farm in eastern Pennsylvania.¹¹⁵ Duck feces on the boots of a farmer may have first brought the virus inside a poultry production facility.¹¹⁶ The virus began benignly, causing drops in egg production and mild upper respiratory symptoms, but soon “rac[ed] though giant commercial chicken warehouses.”¹¹⁷ The now-resident director of the University of Pennsylvania’s poultry laboratory explained that “with that many opportunities to mutate under those intensive conditions,” the virus evolved from being very mild to one described as “chicken Ebola,” causing birds to hemorrhage throughout their bodies.¹¹⁸

Robert Webster’s team performed genetic analyses of the H5N2 virus before and after it turned lethal, and found the two differed by only a single amino acid. The H5 hemagglutinin protein is more than 500 amino acids long.¹¹⁹ A tiny point mutation in the viral genetic material changed the 13th amino acid in the H5 chain from an amino acid named threonine to one called lysine—a mutation that, in Webster’s words, “change[d] that benign virus into one that was completely lethal.”¹²⁰ “That such a tiny change in the virus could enable it to wreak so much havoc,” Webster and colleagues later wrote, “was an awesome discovery.”¹²¹

The U.S. Department of the Interior’s 128-year-old U.S. Geological Survey (USGS), representing the nation’s leading governmental authority on the biological sciences,¹²² reflected on the evolution of low-grade to high-grade strains of bird flu and echoed other world authorities in implicating industrial poultry practices—not only as providing an “excellent opportunity” for rapid spread (particularly when “poultry are housed at high densities in confined quarters”), but, also for playing in part an “ideal” role in possibly sparking the next human pandemic.¹²³

After emerging in 1983, H5N2 resurfaced two years later in low-grade form in Massachusetts, New Jersey, Ohio, and again in Pennsylvania.¹²⁴ Its multi-state resurgence was traced back to live poultry markets in New York City. A survey of live markets in 1986 found 48 harboring the virus,¹²⁵ suggesting to investigators that live poultry markets may have been the critical mixing point between ducks or geese and chickens that triggered the original Pennsylvania epidemic.¹²⁶

Efforts to purge bird flu viruses from live poultry markets over the years have been unsuccessful,¹²⁷ despite periodic quarantine, depopulation, cleaning, and disinfection. In 2004, an outbreak of highly pathogenic H5N2 was discovered in a 7,000-chicken broiler facility in Texas after the owner introduced a chicken from a live poultry market in Houston into his flock.¹²⁸ As described by experts at the USDA and the University of Georgia, “[i]n the United States the live-bird markets that are common in some of the large metropolitan areas can serve as a man-made reservoir of influenza virus....”¹²⁹

H7N2 and Human Infections in the United States

In 2002, the United States saw its first human case of bird flu infection. An H7N2 virus caused more than 200 outbreaks in chicken and turkey operations across a mass poultry production area first in Virginia and then into West Virginia and North Carolina, leading to the destruction of almost 5 million birds.¹³⁰ H7N2 was identified as a low-pathogenicity virus and was suspected in only one human infection,¹³¹ but genetic analyses show that it is mutating toward greater virulence.¹³²

The next year, in 2003, a person was admitted to a hospital in New York with respiratory symptoms and was confirmed to have been infected with the H7N2 bird flu virus. Despite a serious underlying medical condition,

the patient recovered and went home after a few weeks.¹³³ By that year, the virus had swept through millions of egg-laying hens in huge battery-cage facilities in Connecticut and Rhode Island,¹³⁴ outbreaks that the industry admits “confirm the vulnerability of egg production units.”¹³⁵ OIE expert Ilaria Capua describes it as “very difficult” to keep an industrial battery-cage egg facility clean and prevent spread from farm to farm via eggs, egg trays, and equipment. “In our opinion,” she told the Fifth International Symposium on Avian Influenza in 2003, “when the infection gets into a circuit, it will spread within that circuit. It finds its way to spread.”¹³⁶

H7N2’s Spread via Live Bird Markets

The way H7N2 was found to spread in the United States was via live poultry markets. The markets were suspected in the outbreaks in Pennsylvania (1996,¹³⁷ 1997, 1998, 2001, and 2002¹³⁸), Virginia (2002),¹³⁹ Connecticut and Rhode Island (2003), and Delaware (2004).¹⁴⁰ The Delaware outbreak in a broiler chicken operation confining more than 85,000 birds spread to Maryland before it was stopped with the killing of more than 400,000 birds.¹⁴¹ In two cases, direct epidemiological evidence links the presence of trucks hauling birds to live poultry markets at affected farms within one week before the appearance of clinical disease.¹⁴² The trucks deliver birds from the farms to the markets, pick up the empty, dirty crates, and then return to the farms. The “most likely scenario,” according to USDA scientists, is that the crates or trucks were not completely disinfected of the potential billions of infectious particles present with any gross fecal contamination.¹⁴³

The University of Georgia Southeastern Cooperative Wildlife Disease Study describes the U.S. live poultry market system as an “intricate web of retail markets, poultry auctions, wholesale dealers, and farm flocks.” The study notes that in this system, “birds may change hands up to five times before reaching the consumer,” increasing exposure and decreasing trackability.¹⁴⁴

Scientists have watched H7N2 since its emergence in live poultry markets in the United States in 1994. As discussed above, the virus’s fail-safe mechanism that prevents it from becoming too dangerous in its natural waterfowl host is the hemagglutinin activation step that limits viral replication to “safe” organs in the body, like the intestine. However, once placed in a land-based host, such as a chicken, viral mutants that can infect all the victim’s organ systems may have a selective advantage since easy waterborne spread is no longer possible. H5 and H7 viruses can transform by accruing basic (as opposed to acidic or neutral) amino acids in the hemagglutinin protein cleavage site, where the enzymes of the host activate the virus. Once the virus accumulates approximately five basic amino acids, it may transform from a low-pathogenicity virus to one that is highly pathogenic.

The earliest H7N2 isolates in U.S. live poultry markets in 1994 already had two basic amino acids in the critical cleavage site. By 1998, the virus was up to three. In 2002, H7N2 viruses were found with four, needing only one more slight mutation before the virus could have become deadly.¹⁴⁵ Even though the virus was technically still a low-pathogenic strain, the federal government understood the urgency and danger of the virus’s evolution.¹⁴⁶

Ineffective Virus Eradication of Live Bird Markets

Efforts to eradicate the virus at the source—live poultry markets—have failed. According to the USDA Animal and Plant Health Inspection Service, “Despite educational efforts, surveillance, and increased state regulatory efforts, the number of [virus] positive markets has persisted and increased.” In 1998, 30 percent of the markets were infected with H7N2, particularly in the New York metropolitan area. New York has more live markets than all other states in the Northeast combined.¹⁴⁷ By 2001, inspectors could find the virus at 60 percent of markets at any one time.¹⁴⁸

As the states were failing to control the problem and the virus may have needed only one additional mutation before becoming highly pathogenic, the USDA interceded in 2002, coordinating a system-wide closure of all retail live poultry markets throughout the northeastern United States. Following the mass closure, all birds were sold off or killed, and all markets were cleaned and disinfected, left empty for days, and then repopulated with

birds only from closely monitored source flocks confirmed to be negative for all avian influenza viruses. Within five weeks, H7N2 was again found in the live bird markets.

It is unknown whether the virus somehow persisted or was reintroduced. Regardless, despite best efforts at eradication and control, it seems clear that live poultry markets represent a public health risk. Writing in the *Journal of Virology*, USDA researchers concluded that “the rampant reassortment of AIVs [avian influenza viruses] in the LBMs could increase the risk of species crossover because it would increase the chances of the occurrence of the correct constellation of genes to create a virus that replicates efficiently in mammals.”¹⁴⁹

The mass market closure and disinfection did revert the virus’s mutation from four to three basic amino acids.¹⁵⁰ Nevertheless, unless all live poultry markets are closed, H7N2 will presumably continue evolving towards a highly pathogenic strain. According to Cornell University’s Animal Health Diagnostic Center’s virology laboratory director, “It is two major mutations away from becoming a virus that could kill a lot of chickens and become much more pathogenic to people.”¹⁵¹ In 2006, USDA researchers published the results of a study which genetically engineered mutants of an H7N2 virus discovered in New Jersey. They found that the insertion of just a *single* basic amino acid in the right place could transform the virus into a highly pathogenic form.¹⁵² Currently, many suspect that H5N1 is more likely than H7N2 to trigger the next human pandemic, but were it not for H5N1, more attention might be focused on the live poultry markets in New York City—not Hong Kong—to potentially deliver the next avian flu virus with pandemic potential.

Biosecurity: “Wishful Thinking”

Since the primary threat posed by live bird markets is the seeding of industrial poultry operations with virus, the threat could be mediated by meticulous adherence to biosecurity protocols on the farm, such as having all poultry workers, staff, and visitors scrub their footwear in disinfectant every time they stepped into a shed and washing their hands three separate times before entering, as instructed by the USDA’s instructional video *Biosecurity: For the Birds*.¹⁵³ However, as one North Carolina State University poultry health management professor wrote in an industry trade journal, “high biosecurity and proper monitoring are still wishful thinking in many areas of intensive poultry production.”¹⁵⁴

Increasing intensification of animal agriculture industries has made effective biosecurity less of a possibility. The OIE asserts that the changes in the global poultry industry over the past 20 years make infectious diseases “significantly more difficult to control because of the greater number of susceptible animals reared per given unit of time and to the difficulties in applying adequate biosecurity programmes.”¹⁵⁵ The OIE notes that biosecurity measures may be simply incompatible with modern high-density rearing systems.¹⁵⁶ “When an outbreak of avian influenza occurs in an area with a high [poultry] population density,” OIE officials wrote, “the application of rigorous biosecurity measures might not be possible.”¹⁵⁷ The industry and USDA researchers¹⁵⁸ acknowledge that this is one of the disadvantages of the U.S. system: Not only are 20,000 to 25,000 broiler chickens typically confined in a single shed, but there may be up to 16 sheds at a single facility.¹⁵⁹ There are now egg-laying operations with the capacity to cage more than four million hens in a single complex.¹⁶⁰ When sheds and facilities are situated in close enough proximity to one another, once seeded, the virus may be able to spread through the wind.¹⁶¹

Though the U.S. poultry industry claims to have the “best biosecurity system in the world,”¹⁶² academic and governmental investigations have uncovered widespread disregard for biosecurity precautions among both large and small domestic producers.¹⁶³ In 2002, University of Maryland researchers sent questionnaires about biosecurity practices to commercial broiler chicken facilities throughout the Delmarva Peninsula, where more than 100 million chickens raised for meat of various ages are at any given time. Fewer than half the facilities returned the survey, and those that did admitted to severe lapses in biosecurity. The researchers concluded that U.S. broiler chicken flocks “are constantly at risk of infection triggered by poor biosecurity practices.”¹⁶⁴

Charles Beard, acting as U.S. Poultry and Egg Association vice president, admitted that relying on biosecurity measures to protect the U.S. poultry industry “could appear naïve.”¹⁶⁵ “After all, biosecurity is mostly a ‘people’ thing.” Even putting aside human fallibility, Beard is concerned that unless all poultry workers “join in with conviction and enthusiasm, it is not likely to be successful” regardless of what “those in charge” say.¹⁶⁶ The University of Maryland surveys show biosecurity “enthusiasm” to be lacking, and the “convictions” of poultry corporations like Tyson have leaned more toward 20 felony violations for illegal dumping of untreated wastewater into the nation’s rivers¹⁶⁷ than toward a desire to practice biosecurity. In fact, Tyson Foods, the largest chicken-producing corporation in the world,¹⁶⁸ found itself before the Supreme Court in 2005 for refusing to pay workers for time spent donning protective clothing at a poultry plant. The Supreme Court ruled unanimously against Tyson.¹⁶⁹

Breaches in biosecurity occur in modernized facilities around the world.¹⁷⁰ The European Food Safety Authority recognizes that when this happens in a densely populated poultry area, these breaches can result in “massive spread.”¹⁷¹ According to animal disease control experts, biosecurity measures are costly for the industry¹⁷² and “not easy to sustain in the long term.”¹⁷³ Emeritus veterinary poultry professor Simon M. Shane, author of the Handbook on Poultry Diseases, notes a “decline in the standards of biosecurity in an attempt to reduce costs in competitive markets.” The decline is a contributing factor, Shane concludes, in the frequency and severity of disease outbreaks.¹⁷⁴

In *Poultry Digest*, longtime industry and avian health expert Ken Rudd wrote “Poultry Reality Check Needed,” in which he laid out in stark terms the industry’s skewed priorities: “An examination of virtually all the changes made in the past decade shows that they’ve come in the guise of convenience and efficiency, but they are, in fact, cost-cutting measures. Few, if any, decisions have been made solely for the sake of avian health or the long-term protection of the industry. The balance between the two has been lost; the scale is now weighted almost entirely on the cost-cutting side. And, therefore, on the side of microorganisms—much longer on this earth than humans!¹⁷⁵” Specifically, Rudd criticized the profitable yet risky practice of reducing the duration of downtime between flocks and the trend of not cleaning poultry houses between flocks, and pleaded with his industry to “consider the cost of catastrophe.”¹⁷⁶

Examining other widespread viral disease outbreaks, like foot and mouth disease in Europe, the vulnerability of modern animal agriculture is exposed.¹⁷⁷ In the United States, within three years of the emergence of the highly virulent “Delaware variant” of the immunodeficiency infectious bursal disease (IBD) virus, virtually the entire broiler chicken industry east of the Mississippi was affected.¹⁷⁸ The United States, the country that pioneered industrialized poultry production, has reported more bird flu than any other country in the world.¹⁷⁹

The Virginia outbreak in 2002 that led to the deaths of millions of birds and found its way onto hundreds of farms highlights the fallacy of protection in the United States by biosecurity. Based on the rapid domestic spread of bird flu in 2002, leading USDA poultry researchers have concluded the obvious: “[B]iosecurity on many farms is inadequate.”¹⁸⁰ The situation has not necessarily improved since then, according to Yvonne Vizzier Thaxton, executive editor of *Poultry* magazine and professor of poultry science at Mississippi State. In 2005, she editorialized that “I believe it is time to reexamine biosecurity in our industry. We’ve become lax in many ways, and this is exactly what it took to get the 1983 AI outbreak moving...If WHO is right and a pandemic brings human AI to the United States, will you be able to look your family and neighbors in the eye and say you’ve done all you can to stop the spread? Having to answer that question alarms me!”¹⁸¹

Closing Down Live Bird Markets in Asia

H5N1 is considered a Biosafety Level 3+ pathogen,¹⁸² meaning that, in a laboratory setting, the virus is only to be handled in unique, high-containment buildings specially engineered with air locks, controlled access corridors, and double-door entries. Access is limited to competent personnel with extensive training,¹⁸³ and showering is required upon each entry and exit.¹⁸⁴ Air flow is ducted for unidirectional, single-pass filtered exhaust only.¹⁸⁵ All floors, walls, and ceilings are waterproofed and sealed with continuous cove moldings.¹⁸⁶ All wall penetrations,

such as electrical outlets and phone lines, are caulked, collared, or sealed to prevent any leaks.¹⁸⁷ Surfaces are disinfected on a daily basis,¹⁸⁸ and solid wastes are incinerated.¹⁸⁹ “Unfortunately,” leading USDA poultry virologist Dennis Senne told an international gathering of bird flu scientists, “that level of biosecurity does not exist in U.S. poultry production and I doubt that it exists in other parts of the world.”¹⁹⁰

Recognizing both the seriousness of the public health threat and the impracticality of absolute biosecurity, governments throughout Asia are phasing out these markets. China has set the precedent of attempting¹⁹¹ to ban all live bird markets in Shanghai, its largest city, as well as the capital city of Beijing.¹⁹² Hong Kong has also decided to phase them out,¹⁹³ and the Chinese government has reportedly urged all large cities to “gradually call off the killing and sales of live fowls in the market.”¹⁹⁴ Similarly, bans have reportedly been proposed in Singapore, Japan,¹⁹⁵ and Vietnam’s Hanoi, Hai Phong, Vinh and Ho Chi Minh City.¹⁹⁶ Starting in 2008, those found publicly slaughtering birds in Taiwan may face a 500,000 NT fine (approximately \$15,000 USD).¹⁹⁷ Though there has never been a recorded outbreak in Taiwan, the chairman of its National Science Council explained the reasoning behind the ban: “We can’t foresee whether an outbreak of bird flu will happen in Taiwan, but every nation in the world is obligated to take part in the prevention of the epidemic.”¹⁹⁸

There has been support from the scientific community for the transition from live markets to supermarkets. Paul Chan, for example, a professor of microbiology at Chinese University of Hong Kong, told the *China Daily*, “I support doing away with selling live poultry altogether.”¹⁹⁹ In light of the emergence of SARS and H5N1 from live animal markets in Asia, a 2006 scientific review concluded that “the most optimal strategy is to forbid all kinds of live animals at wet-markets, with enforcement of central slaughtering.”²⁰⁰

Robert Webster, the world’s authority on avian influenza and chair of the Virology Division of St. Jude Children’s Research Hospital in Tennessee and director of the U.S. Collaborating Center of the WHO, believes that even with all Hong Kong live markets closed, the overall pandemic risk may not be sufficiently affected unless live markets could be closed throughout China and elsewhere.²⁰¹ Webster’s team has concluded, “Until the traditional practice of selling poultry in the live market changes we will have to accept that live markets are breeding grounds for influenza viruses....”²⁰²

Live Bird Markets Should Be Shut Down Throughout the United States

Though suspected of playing a pivotal role in the spread of bird flu in the United States,²⁰³ live bird markets in this country are proliferating. In New York City, for example, the number of live bird markets almost doubled from 44 in 1994 to more than 80 in 2002.²⁰⁴ Given the risk, many in the U.S. commercial poultry industry are “absolutely determined” to have live markets eliminated, according to a Louisiana State University poultry scientist.²⁰⁵ Briefly but incisively, the president of the USA Poultry and Egg Export Council commented, “We can’t jeopardize the entire U.S. [poultry] industry.”²⁰⁶

According to the USDA’s Agricultural Research Service, “The U.S. currently has the largest, most genetically homogeneous and, thus potentially, the most disease-susceptible population of food animals in the history of mankind....The emergence of a new disease or a slight shift in the epidemiology of an existing disease could lead to immediate and disastrous results for American livestock producers and consumers.”²⁰⁷

The USDA agrees that live bird markets have been shown to present a “major risk” to the nation’s poultry industry.²⁰⁸ Agency scientists wrote that “live bird markets of the Northeast remain the biggest concern for the presence of avian influenza in the United States.”²⁰⁹ Yet, despite agreement by both the industry and the USDA, live bird markets persist and indeed flourish. Some industry officials fear that closing bird markets would drive the entire trade underground, making it even more difficult to regulate.²¹⁰ After the SARS outbreak, for example, customer demand in Asia drove the cost of civet cats up to \$200, making it likely that such animals could be obtained regardless of legality.²¹¹ The USDA has therefore chosen to “manage or mitigate the risk rather than to outlaw it.”²¹²

By its own admission, the USDA is doing a poor job of risk management. Speaking at the Fifth International Symposium on Avian Influenza in 2002, USDA poultry researchers said, “Considerable efforts are continuing on the part of industry and state and federal governments to control influenza in the LBM system, but currently the efforts have been unsuccessful.”²¹³ “Despite educational efforts, surveillance, and increased state regulatory efforts,” the USDA admitted the following year that “the number of [bird flu] positive markets has persisted and increased.”²¹⁴ Live bird markets seem inherently risky. In response to virus isolations from New Jersey’s markets, the state veterinarian said, “They can be doing everything right and still have a market that tests positive.”²¹⁵

Even if a segment of the live bird trade was forced underground, the situation and risks might not significantly worsen. Record-keeping in live poultry markets is already sparse or nonexistent even as to the birds’ country of origin.²¹⁶ Currently, the purchase and sale of live birds is a cash business in which market owners are “disinclined to keep accurate records that would be costly if subjected to IRS scrutiny.”²¹⁷ A 2003 survey of handling practices at live bird markets found that fewer than 2 percent of suppliers followed the recommended biosecurity practices to prevent the spread of the disease.²¹⁸ USDA Science Hall of Famer²¹⁹ Charles Beard is concerned that U.S. live bird markets could be the portal by which H5N1 enters commercial poultry flocks in the United States.²²⁰

Live bird markets continue to exist in the United States only because local health authorities continue to license them. They are exempt from federal meat inspection laws because they slaughter fewer than 20,000 birds a year,²²¹ an exemption that doesn’t apply for other animals.²²² Poultry specialists predict that if live bird markets had to be held to the same federal standards of inspection, cleanliness, and pathogen control as the commercial poultry industry, or small producers of other animals, “authorities could virtually eliminate LBMs.”²²³

Currently, all regulation of live markets, including licensing and inspection, is done at a state level and, as such, represents a patchwork with some states having little or no intact regulatory apparatus. To remedy the situation, in October 2004 the USDA released a guidance document with uniform standards for regulation, suggesting quarterly testing and temporary market closure and disinfection if H5 and H7 AI viruses are discovered. By 2006, the USDA asserted they had Memorandums of Understanding from all high-risk states, promising to move toward these standardized guidelines. In exchange, the states receive federal funds, resources, and assistance from staff, if needed, to carry out their oversight of the markets. As a result of increased surveillance, the number of markets found to be positive for AI viruses is reportedly in decline.²²⁴ However, despite these efforts, experts at the USDA and the University of Georgia concluded in a 2006 review that U.S. live bird markets remain “an ideal environment for transmission, adaptation and evolution of avian influenza viruses.”²²⁵

The regulatory strategy of increased surveillance and disinfection was developed at a time when avian influenza was only known to affect birds. With knowledge that these viruses can kill people directly, can escape into the environment and, in fact, were the source of the 1918 pandemic—the worst plague in human history—the United States must follow Asia’s example and shut down these markets.

The authority conferred upon the Secretary of Health and Human Services by Title 42 United States Code Section 264 to prevent the introduction, transmission, and spread of communicable diseases in the United States could be used to close live bird markets across the country. Just as the Food and Drug Administration banned the sale of small turtles for pets in 1975 to prevent the spread of *Salmonella*, it (or another appropriate agency, such as the Centers for Disease Control and Prevention) should be able to ban the sale of birds at live bird markets to prevent the spread of avian influenza.

On January 3, 2007, New York City Assemblywoman Barbara Clark introduced NY A.B. 246, which would prohibit the licensing of new establishments that slaughter animals within 25 feet of a residential dwelling in a city with a population of one million or more. This would effectively outlaw any new live bird markets in New York City, where markets and their public health risks remain most concentrated.

Conclusion

In a 2004 consultation document for the government, the University of Hong Kong School of Public Health explained what was at stake: “The trade-off between the preference for eating the flesh of freshly slaughtered chicken and the risk to local, regional, and global population health from avian influenza should be addressed directly, and in terms of a precautionary public health approach aimed at providing the greatest benefit to the maximum possible number of people.”²²⁶ The United States government should follow suit and find that the risks of live bird markets far outweigh justifications for their existence.

Dr. Webster concluded a landmark article on the emergence of pandemic strains of influenza with these words: “An immediate practical approach is to close all live poultry markets....” He notes that with refrigeration systems widely available—even through much of the developing world—it is no longer necessary to sell live birds. “The reality is that traditions change very slowly,” he said, but “a new pandemic could accelerate this process.”²²⁷ There is a tendency not to “shore up the levies” until after disaster strikes, but by acting now to close live bird markets in the United States, the global risk of the future emergence of flu viruses with pandemic potential may be reduced.

References

1. Senne DA, Pearson JE, and Panigrahy B. 1997. Live poultry markets: a missing link in the epidemiology of avian influenza. In: Proceedings of the 3rd International Symposium on Avian Influenza, May 27-29 (University of Wisconsin, Madison, pp. 50-8).
2. Webster RG. 1997. Influenza virus: transmission between species and relevance to emergence of the next human pandemic. *Archives of Virology* 13:S105-13.
3. Yousaf M. 2004. Avian influenza outbreak hits the industry again. *World Poultry* 20(3):22-5.
4. Webster RG, Bean WJ, Gorman OT, Chambers TM, and Kawaoka Y. 1992. Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56(1):152-79.
5. Webster RG, Bean WJ, Gorman OT, Chambers TM, and Kawaoka Y. 1992. Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56(1):152-79.
6. Laver WG, Bischofberger N, and Webster RG. 2000. The origin and control of pandemic influenza. *Perspectives in Biology and Medicine* 43(2):173-92.
7. Wobeser GA. 1997. *Diseases of Wild Waterfowl* (New York, NY: Plenum Press).
8. Webster RG, Shortridge KF, and Kawaoka Y. 1997. Influenza: interspecies transmission and emergence of new pandemics. *Federation of European Microbiological Societies Immunology and Medical Microbiology* 18:275-9.
9. Murphy B. 1993. Factors restraining emergence of new influenza viruses. In: Morse SS (ed.), *Emerging Viruses* (New York, NY: Oxford University Press, pp. 234-40).
10. Drexler M. 2002. *Secret Agents: The Menace of Emerging Infections* (Washington, DC: Joseph Henry Press).
11. Webster RG, Wright SM, Castrucci MR, Bean WJ, and Kawaoka Y. 1993. Influenza: a model of an emerging virus disease. *Intervirology* 35:16-25.
12. Shortridge KF. 1992. Pandemic influenza: a zoonosis? *Seminars in Respiratory Infections* 7:11-25.
13. Webster RG, Bean WJ, Gorman OT, Chambers TM, and Kawaoka Y. 1992. Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56(1):152-79.
14. Webby RJ and Webster RG. 2001. Emergence of influenza A viruses. *Philosophical Transactions of the Royal Society of London* 356:1817-28.
15. Webster RG, Bean WJ, Gorman OT, Chambers TM, and Kawaoka Y. 1992. Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56(1):152-79.
16. Handwerk B. 2002. Sharks falling prey to humans' appetites. *National Geographic News*, June 3. news.nationalgeographic.com/news/2002/06/0603_020603_shark1.html. Accessed April 1, 2007.
17. Sharp GB, Kawaoka Y, Jones DJ, et al. 1997. Coinfection of wild ducks by influenza A viruses: distribution patterns and biological significance. *Journal of Virology* 71:6128-35. jvi.asm.org/cgi/reprint/71/8/

- 6128.pdf. Accessed April 1, 2007.
18. Kurtz J, Manvell RJ, and Banks J. 1996. Avian influenza virus isolated from a woman with conjunctivitis. *Lancet*. 348(9031):901-2.
 19. Webster RG, Geraci J, Petursson G, and Skirnisson K. 1981. Conjunctivitis in human beings caused by influenza A virus of seals. *New England Journal of Medicine* 304(15):911.
 20. Davis M. 2005. *The Monster at Our Door: The Global Threat of Avian Flu* (New York, NY: The New Press).
 21. Subbarao K and Katz J. 2000. Avian influenza viruses infecting humans. *Cellular and Molecular Life Sciences* 57:1770-84.
 22. Webby RJ and Webster RG. 2001. Emergence of influenza A viruses. *Philosophical Transactions of the Royal Society of London* 356:1817-28.
 23. Whitney D. 2005. Bird flu battle on horizon: scientists fear the deadly virus is adapting and will hit the U.S. hard. *Sacramento Bee*, August 13, p. A3.
 24. Honigsbaum M. 2005. Flying Dutchman to the rescue: "virus hunter" sees bird flu as greatest threat. *Guardian*, June 3, p. 23. www.guardian.co.uk/life/feature/story/0,13026,1491811,00.html. Accessed April 1, 2007.
 25. World Health Organization. 2005. Avian influenza: assessing the pandemic threat, January 1. www.who.int/csr/disease/influenza/H5N1-9reduit.pdf. Accessed April 1, 2007.
 26. Suarez DL, Garcia M, Latimer J, Senne D, and Perdue M. 1999. Phylogenetic analysis of H7 avian influenza viruses isolated from the live bird markets of the northeastern United States. *Journal of Virology* 73:3567-73.
 27. Webster RG. 1998. Influenza: an emerging microbial pathogen. In: *Emerging Infections* (San Diego, CA: Academic Press, pp. 275-300).
 28. Dronamraju K (ed.). 2004. *Infectious Disease and Host-Pathogen Evolution* (Cambridge, UK: Cambridge University Press).
 29. Hollenbeck JE. 2005. An avian connection as a catalyst to the 1918-1919 influenza pandemic. *International Journal of Medical Sciences* 2(2):87-90. www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1145139. Accessed April 1, 2007.
 30. Suarez DL. 2000. Evolution of avian influenza viruses. *Veterinary Microbiology* 74:15-27.
 31. Suarez DL, Perdue ML, Cox N, et al. 1998. Comparisons of highly virulent H5N1 influenza A viruses isolated from humans and chickens from Hong Kong. *Journal of Virology* 72:6678-88.
 32. Shortridge KF. 1992. Pandemic influenza: a zoonosis? *Seminars in Respiratory Infections* 7:11-25.
 33. Drexler M. 2002. *Secret Agents: The Menace of Emerging Infections* (Washington, DC: Joseph Henry Press).
 34. Capua I and Alexander DJ. 2004. Avian influenza: recent developments. *Avian Pathology* 33:393-404.
 35. Sturm-Ramirez KM, Hulse-Post DJ, and Govorkova EA. 2005. Are ducks contributing to the endemicity of highly pathogenic H5N1 influenza virus in Asia? *Journal of Virology* 79(17):11269-79.
 36. Marwick C. 1998. Investigators present latest findings on Hong Kong "bird flu" to the FDA. *Journal of the American Medical Association* 279(9):643-4.
 37. Van Blerkom LM. 2003. Role of viruses in human evolution. *Yearbook of Physical Anthropology* 46:14-46.
 38. Dimmock NJ, Easton A, and Leppard K. 2001. *Introduction to Modern Virology* (Oxford, UK: Blackwell Publishing).
 39. Davis M. 2005. *The Monster at Our Door: The Global Threat of Avian Flu* (New York, NY: The New Press).
 40. Suarez DL. 2000. Evolution of avian influenza viruses. *Veterinary Microbiology* 74:15-27.
 41. Ito T, Goto H, Yamamoto E, et al. Generation of a highly pathogenic avian influenza A virus from an avirulent field isolate by passaging in chickens. *Journal of Virology* 75(9):4439-43. www.pubmedcentral.com/articlerender.fcgi?artid=114193. Accessed April 1, 2007.
 42. Capua I and Marangon S. 2003. The use of vaccination as an option for the control of avian influenza. In: *71st General Session International Committee of the World Organization for Animal Health* (Paris, France, May 18-23).

43. Morris RS and Jackson R. 2005. Epidemiology of H5N1 avian influenza in Asia and implications for regional control. Food and Agriculture Organization of the United Nations. January-February 11. www.thepoultrysite.com/FeaturedArticle/FAType.asp?AREA=turkeys&Display=121. Accessed April 1, 2007.
44. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
45. U.S. Department of Agriculture Agriculture Research Service. 2006. People and places. www.ars.usda.gov/pandp/people/people.htm?personid=5507. Accessed April 1, 2007.
46. Ito T, Goto H, Yamamoto E, et al. Generation of a highly pathogenic avian influenza A virus from an avirulent field isolate by passaging in chickens. *Journal of Virology* 75(9):4439-43. www.pubmedcentral.com/articlerender.fcgi?artid=114193. Accessed April 1, 2007.
47. Stegeman A. 2003. Workshop 1: introduction and spread of avian influenza. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control*. library.wur.nl/frontis/avian_influenza/workshop1.pdf. Accessed April 1, 2007.
48. For more information, see: An HSUS Report: Human Health Implications of Intensive Poultry Production and Avian Influenza at www.hsus.org/farm/resources/research/pubhealth/public_health_avian_influenza.html. Accessed March 25, 2007.
49. Ritchie BW. 1995. *Avian Viruses: Function and Control* (Lake Worth, FL: Wingers Publishing).
50. Delgado C, Rosegrant M, Steinfeld H, Ehui S, and Courbois C. 1999. *Livestock to 2020: the next food revolution*. Food, Agriculture, and the Environment Discussion Paper 28. For the International Food Policy Research Institute, the Food and Agriculture Organization of the United Nations and the International Livestock Research Institute. www.ifpri.org/2020/dp/dp28.pdf. Accessed February 2, 2007.
51. Hafez HM. 2003. Emerging and re-emerging diseases in poultry. *World Poultry* 19(7):23-7.
52. Ritchie BW, op. cit.
53. Frederick MA. 1999. The threat posed by the global emergence of livestock, food-borne, and zoonotic pathogens. *Annals of the New York Academy of Sciences* 894:20-7.
54. André F and Tuytens M. 2005. The importance of straw for pig and cattle welfare: a review. *Applied Animal Behavior Science* 92(3):261.
55. Maes D, Deluyker H, Verdonck M, et al. 2000. Herd factors associated with the seroprevalences of four major respiratory pathogens in slaughter pigs from farrow-to-finish pig herds. *Veterinary Research* 31:313-27.
56. Quattro JD. 1999. Three scientists introduced into ARS hall of fame. *ARS/USDA News and Events*, September 17. www.ars.usda.gov/is/pr/1999/990917.htm. Accessed February 2, 2007.
57. Witter RL. 1998. Control strategies for Marek's Disease: a perspective for the future. *Poultry Science* 77:1197-203.
58. El-Lethey H, Huber-Eicher B, and Jungi TW. 2003. Exploration of stress-induced immunosuppression in chickens reveals both stress-resistant and stress-susceptible antigen responses. *Veterinary Immunology and Immunopathology* 95:91-101.
59. For more information, see: An HSUS Report: Welfare Issues with Selective Breeding Rapid Growth in Broiler Chickens and Turkeys at www.hsus.org/farm/resources/research/practices/fast_growth_broilers.html. Accessed March 26, 2007.
60. Wilson WO. 1966. Poultry production. *Scientific American* 215(1):56-64.
61. U.S. Department of Agriculture Animal and Plant Health Inspection Service. 2004. Highly pathogenic avian influenza: a threat to U.S. poultry. Program Aid No. 1704. March.
62. Cole DJ, Hill VR, Humenik FJ, and Sobsey MD. 1999. Health, safety, and environmental concerns of farm animal waste. *Occupational Medicine: State of the Art Reviews* 14(2):423-48.
63. North MO and Bell DD. 1990. *Commercial Chicken Production Manual*, 4th Edition (New York, NY: Van Nostrand Reinhold).
64. European Commission Scientific Committee on Animal Health and Animal Welfare. 2000. *The Welfare of Chickens Kept for Meat Production (Broilers)*. March 21. europa.eu.int/comm/food/fs/sc/scah/out39_en.pdf. Accessed April 2, 2007.

65. Madec F and Rose N. 2003. How husbandry practices may contribute to the course of infectious diseases in pigs. In: 4th International Symposium on Emerging and Re-emerging Pig Diseases (Rome, Italy June 29th-July 2nd, pp. 9-18).
66. Cooper GL, Venables LM, and Lever MS. 1996. Airborne challenge of chickens vaccinated orally with the genetically-defined *Salmonella enteritidis* aroA strain CVL30. *Veterinary Record* 139(18):447-8.
67. Van der Sluis W. 2005. Housing conditions affect broiler welfare more than stocking density. *World Poultry* 21(8):22-3.
68. Stohr K and Meslin FX. 1997. The role of veterinary public health in the prevention of zoonoses. *Archives Virology* 13:S207-18.
69. Pheasant B. 2004. A virus of our hatching. *The Financial Review* (Australia), January 30.
70. Capua I and Marangon S. 2003. The use of vaccination as an option for the control of avian influenza. World Organization for Animal Health 71st General Session in Paris, France, May 18-23. www.oie.int/eng/avian_influenza/a_71%20sg_12_cs3e.pdf. Accessed April 1, 2007.
71. Vidal J. 2006. Flying in the face of nature. *Guardian*, February 22. society.guardian.co.uk/health/story/0,,1714703,00.html. Accessed April 1, 2007.
72. Zhihong H, Shuyi Z, and Zhu C. 2005. Zoonoses: the deadly diseases that animals pass on to us. *Global Agenda: Magazine of World Economic Forum Annual Meeting*.
73. Pheasant B. 2004. A virus of our hatching. *The Financial Review* (Australia), January 30.
74. Chastel C. 2004. Emergence of new viruses in Asia: is climate change involved? *Médecine et maladies infectieuses* 34:499-505.
75. Prowse S. 2005. Biosecurity and emerging infectious diseases. *Australia Academy of Technological Sciences and Engineering Focus* No 136, March/April.
76. Boyd W. 2001. Making meat: science, technology, and American poultry production. *Technology and Culture* 42:631-64.
77. United Nations. 2005. UN task forces battle misconceptions of avian flu, mount Indonesian campaign. UN News Centre, October 24. www.un.org/apps/news/story.asp?NewsID=16342&Cr=bird&Cr1=flu. Accessed April 1, 2007.
78. United Nations. 2005. UN task forces battle misconceptions of avian flu, mount Indonesian campaign. UN News Centre, October 24. www.un.org/apps/news/story.asp?NewsID=16342&Cr=bird&Cr1=flu. Accessed April 1, 2007.
79. 2005. Preeminent vaccinologist led eradication of world's deadliest childhood diseases, saving and protecting millions. *American Association of Immunologists Newsletter*, June/July, p. 8. www.aai.org/newsletter/PDFs/2005/JunJulPDF.pdf. Accessed April 7, 2007.
80. Prince Mahidol Award Foundation. 1998. Professor Kennedy F Shortridge, Ph.D. www.kanchanapisek.or.th/pmaf/laureate-bio.en.php?type=ind&id=2005-12-26%2008:58:08. Accessed April 7, 2007.
81. 2004. Avian influenza researcher to receive UQ honorary doctorate. University of Queensland. September 8. www.uq.edu.au/news/?article=5877. Accessed April 7, 2007.
82. Cooper S. 2006. (How to stop) The next killer flu. *Seed Magazine*, February-March.
83. Torrey EF and Yolken RH. 2005. *Beasts of the Earth: Animals, Humans, and Disease* (New Brunswick, NJ: Rutgers University Press).
84. Webby RJ and Webster RG. 2001. Emergence of influenza A viruses. *Philosophical Transactions of the Royal Society of London* 356:1817-28.
85. Shortridge KF, Peiris JS, and Guan Y. 2003. The next influenza pandemic: lessons from Hong Kong. *Journal of Applied Microbiology* 94(1):70.
86. Brown C. 2004. Emerging zoonoses and pathogens of public health significance: an overview. *Revue Scientifique et Technique* 23:435-42.
87. Fielding R, Lam WW, Ho EY, Lam TH, Hedley AJ, and Leung GM. 2005. Avian influenza risk perception, Hong Kong. *Emerging Infectious Diseases* 11(5):677-82.
88. Senne DA, Pederson JC, and Panigrahy B. 2003. Live-bird markets in the northeastern United States: a source of avian influenza in commercial poultry. In: Schrijver RS and Koch G (eds.), *Proceedings of the*

- Frontis Workshop on Avian Influenza: Prevention and Control (Wageningen, The Netherlands, pp. 19-24).
89. Parry J. 2003. Hong Kong under WHO spotlight after flu outbreak. *British Medical Journal* 327:308.
 90. Fong J. 2004. Beating “bird flu” against the odds. *Greatreporter.com*, March 3. greatreporter.com/mambo/content/view/227/2/. Accessed April 1, 2007.
 91. Macan-Markar M. 2005. Bird flu pandemic may hang on for years. *Inter Press Service*, February 26.
 92. Dronamraju K (ed.). 2004. *Infectious Disease and Host-Pathogen Evolution* (Cambridge, UK: Cambridge University Press).
 93. Kingsbury K. 2004. Animal markets are alive with controversy. *Columbia News Service*. February 16. www.jrn.columbia.edu/studentwork/cns/2004-02-16/455.asp. Accessed April 1, 2007.
 94. Appenzeller T. 2005. Tracking the next killer flu. *National Geographic*, October. www7.nationalgeographic.com/ngm/0510/feature1/. Accessed April 1, 2007.
 95. Webster RG and Hulse DJ. 2004. Microbial adaptation and change: avian influenza. *Revue Scientifique et Technique* 23(2):453-65.
 96. Stubbs EL. 1948. Fowl pest. In: Biester HE and Schwarte LH (eds.), *Diseases of Poultry*, 2nd ed (Ames, IA: Iowa State University Press, pp. 603-14).
 97. Perez DR, Nazarian SH, McFadden G, and Gilmore MS. 2005. Miscellaneous threats: highly pathogenic avian influenza, and novel bio-engineered organisms. In: Bronze MS and Greenfield RA (eds.), *Biodefense: Principles and Pathogens* (Norfolk, UK: Horizon Bioscience).
 98. Senne DA, Pearson JE, and Panigrahy B. 1997. Live poultry markets: a missing link in the epidemiology of avian influenza. In: *Proceedings of the 3rd International Symposium on Avian Influenza May 27-29* (University of Wisconsin, Madison, pp. 50-8).
 99. Perez DR, Nazarian SH, McFadden G, and Gilmore MS. 2005. Miscellaneous threats: highly pathogenic avian influenza, and novel bio-engineered organisms. In: Bronze MS and Greenfield RA (eds.), *Biodefense: Principles and Pathogens* (Norfolk, UK: Horizon Bioscience).
 100. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. www.findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
 101. Personal communication with Dennis Thompson, Chief of the Meat and Poultry Inspection Branch of the California Department of Food and Agriculture, July 6, 2006.
 102. Personal communication with Dale Lauer, Director of Poultry Programs, Minnesota Poultry Testing Laboratory. Minnesota Board of Animal Health, July 11, 2006.
 103. Personal communication with Howard C. “Butch” Johnson, State Director and Manager, Meat Safety Assurance of the Texas Department of State Health Services, July 21, 2006.
 104. Personal communication with Jennifer Glover, poultry specialist in the Animal Disease Control Section of the Florida Department of Agriculture and Consumer Services, July 11, 2006.
 105. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
 106. Halvorson D. 2005. Overview of avian influenza. University of Minnesota University of Minnesota Extension Service. December 9. www.cvm.umn.edu/ai/home.html. Accessed April 1, 2007.
 107. U.S. Department of Agriculture Animal and Plant Inspection Service. Avian influenza in the United States. oars.aphis.usda.gov/lpa/issues/ai_us/ai_us.html. Accessed April 1, 2007.
 108. Brown C. 1999. Economic considerations of agricultural diseases. *Annals of the New York Academy of Sciences* 894:92-4.
 109. Swayne DE and Akey BL. 2003. Avian influenza control strategies in the United States of America. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 113-30).
 110. Brown C. 1999. Economic considerations of agricultural diseases. *Annals of the New York Academy of Sciences* 894:92-4.
 111. Alexander DJ. 2000. A review of avian influenza in different bird species. *Veterinary Microbiology* 74:3-13.

112. Stubbs EL. 1925. Fowl pest. Sixty-Second Annual Meeting of the American Veterinary Medical Association, Portland, Oregon, July 21-24.
113. Linares JA, Gayle L, Sneed L, and Wigle W. 2004. H5N2 avian influenza outbreak in Texas. In: 76th Northeastern Conference on Avian Diseases: June 9-11 (State College, Pennsylvania: Department of Veterinary Science, College of Agricultural Sciences, Pennsylvania State University, p. 14). www.vetsci.psu.edu/NECAD/NECADProceedings.pdf. Accessed April 1, 2007.
114. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
115. Senne DA, Pearson JE, and Panigrahy B. 1997. Live poultry markets: a missing link in the epidemiology of avian influenza. In: *Proceedings of the 3rd International Symposium on Avian Influenza May 27-29* (University of Wisconsin, Madison, pp. 50-8).
116. Gladwell M. 1995. The plague year. *New Republic*, July 24.
117. Laver WG, Bischofberger N, and Webster RG. 2000. The origin and control of pandemic influenza. *Perspectives in Biology and Medicine* 43(2):173-92.
118. Gladwell M. 1997. The dead zone. *New Yorker*, September 9. www.gladwell.com/1997/1997_09_29_a_flu.htm. Accessed April 7, 2007.
119. Branden C and Tooze J. 1991. *Introduction to Protein Structure* (New York, NY: Garland Publishing, Inc., pp. 72-5).
120. Kawaoka Y and Webster RG. 1988. Molecular mechanism of acquisition of virulence in influenza virus in nature. *Microbial Pathogenesis* 5:311-8.
121. Laver WG, Bischofberger N, and Webster RG. 2000. The origin and control of pandemic influenza. *Perspectives in Biology and Medicine* 43(2):173-92.
122. Dierauf L. Avian influenza in wild birds. U.S. Department of the Interior U.S. Geological Survey. *Wildlife Health Bulletin* 04-01. www.nwhc.usgs.gov/publications/wildlife_health_bulletins/WHB_04_01.jsp. Accessed April 1, 2007.
123. Dierauf L. Avian influenza in wild birds. U.S. Department of the Interior U.S. Geological Survey. *Wildlife Health Bulletin* 04-01. www.nwhc.usgs.gov/publications/wildlife_health_bulletins/WHB_04_01.jsp. Accessed April 1, 2007.
124. Senne DA, Pearson JE, and Panigrahy B. 1997. Live poultry markets: a missing link in the epidemiology of avian influenza. In: *Proceedings of the 3rd International Symposium on Avian Influenza May 27-29* (University of Wisconsin, Madison, pp. 50-8).
125. Senne DA, Pearson JE, and Panigrahy B. 1997. Live poultry markets: a missing link in the epidemiology of avian influenza. In: *Proceedings of the 3rd International Symposium on Avian Influenza May 27-29* (University of Wisconsin, Madison, pp. 50-8).
126. Perez DR, Nazarian SH, McFadden G, and Gilmore MS. 2005. Miscellaneous threats: highly pathogenic avian influenza, and novel bio-engineered organisms. In: *Bronze MS and Greenfield RA (eds.), Biodefense: Principles and Pathogens* (Norfolk, UK: Horizon Bioscience).
127. Lee C, Senne DA, Linares JA, et al. 2004. Characterization of recent H5 subtype avian influenza viruses from U.S. poultry. *Avian Pathology* 33:288-97.
128. 2004. USDA confirms highly pathogenic avian influenza in Texas. *West Texas County Courier*, February 26. www.wtccourier.com/flats_pdf/2004/02-26-04.pdf. Accessed April 7, 2007.
129. Senne DA, Suarez DL, Stallnecht E, Pedersen JC, and Panigrahy B. 2006. Ecology and epidemiology of avian influenza in North and South America. *Developments in Biologicals* 124:37-44.
130. Senne DA, Holt TJ, and Akey BL. 2003. An overview of the 2002 outbreak of low-pathogenic H7N2 avian influenza in Virginia, West Virginia and North Carolina. In: *Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 41-7).
131. Centers for Disease Control and Prevention. 2006. Avian Influenza Infection in Humans. www.cdc.gov/flu/avian/gen-info/avian-flu-humans.htm. Accessed April 1, 2007.
132. Spackman E and Suarez DL. 2003. Evaluation of molecular markers for pathogenicity in recent H7N2 avian influenza isolates from the northeastern United States. In: *Proceedings of the 52nd Western Poultry Disease Conference*, pp. 21-3 (Sacramento, CA).

133. Centers for Disease Control and Prevention. 2006. Past Avian Influenza Outbreaks. www.cdc.gov/flu/avian/outbreaks/past.htm. Accessed April 7, 2007.
134. Swayne DE and Akey BL. 2003. Avian influenza control strategies in the United States of America. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 113-30).
135. Shane SM. 2003. Disease continues to impact the world's poultry industries. *World Poultry* 19(7):22-7.
136. Swayne DE. 2003. Transcript of the question and answer session from the Fifth International Symposium on Avian Influenza. *Avian Diseases* 47:1219-55.
137. Senne DA, Pederson JC, and Panigrahy B. 2003. Live-bird markets in the northeastern United States: a source of avian influenza in commercial poultry. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 19-24).
138. Spackman E, Senne DA, Davison S, and Suarez DL. 2003. Sequence analysis of recent H7 avian influenza viruses associated with three different outbreaks in commercial poultry in the United States. *Journal of Virology* 77:13399-402.
139. Swayne DE and Akey BL. 2003. Avian influenza control strategies in the United States of America. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 113-30).
140. Centers for Disease Control and Prevention. 2006. Past Avian Influenza Outbreaks. www.cdc.gov/flu/avian/outbreaks/past.htm. Accessed April 7, 2007.
141. Capua I and Alexander DJ. 2004. Avian influenza: recent developments. *Avian Pathology* 33:393-404.
142. Senne DA, Pederson JC, and Panigrahy B. 2003. Live-bird markets in the northeastern United States: a source of avian influenza in commercial poultry. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 19-24).
143. Swayne DE. 2003. Transcript of the question and answer session from the Fifth International Symposium on Avian Influenza. *Avian Diseases* 47:1219-55.
144. University of Georgia Southeastern Cooperative Wildlife Disease Study. 1986. July. Avian influenza in live poultry markets. *SCWDS Briefs*, July.
145. Spackman E, Senne DA, Davison S, and Suarez DL. 2003. Sequence analysis of recent H7 avian influenza viruses associated with three different outbreaks in commercial poultry in the United States. *Journal of Virology* 77:13399-402.
146. Swayne DE and Akey BL. 2003. Avian influenza control strategies in the United States of America. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 113-30).
147. Mullaney R. 2003. Live-bird market closure activities in the northeastern United States. *Avian Diseases* 47:1096-8.
148. Senne DA, Pederson JC, and Panigrahy B. 2003. Live-bird markets in the northeastern United States: a source of avian influenza in commercial poultry. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 19-24).
149. Suarez DL, Garcia M, Latimer J, Senne D, and Perdue M. 1999. Phylogenetic analysis of H7 avian influenza viruses isolated from the live bird markets of the northeastern United States. *Journal of Virology* 73:3567-73.
150. Senne DA, Pederson JC, and Panigrahy B. 2003. Live-bird markets in the northeastern United States: a source of avian influenza in commercial poultry. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands, pp. 19-24).
151. 2005. Cornell checking for avian flu in NYC. *Ithaca Journal*, October 15.
152. Lee CW, Lee YJ, Senne DA, and Suarez DL. 2006. Pathogenic potential of North American H7N2 avian influenza virus: a mutagenesis study using reverse genetics. *Virology* 353(2):388-95.
153. U.S. Department of Agriculture Animal and Plant Health Inspection Service. Backyard biosecurity: practices to keep your birds healthy. September 2004
154. Vaillancourt JP. 2002. Biosecurity now. *Poultry International*. 41(8):12-8.

155. Capua I and Marangon S. 2003. The use of vaccination as an option for the control of avian influenza. In: 71st General Session International Committee of the World Organization for Animal Health (Paris, France, May 18-23).
156. Capua I and Marangon S. 2003. The use of vaccination as an option for the control of avian influenza. In: 71st General Session International Committee of the World Organization for Animal Health (Paris, France, May 18-23).
157. Capua I and Marangon S. 2003. Currently available tools and strategies for emergency vaccination in case of avian influenza. In: Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control (Wageningen, The Netherlands, pp. 59-74).
158. Perdue ML, Suarez DL, and Swayne DE. 2000. Avian influenza in the 1990s. *Avian and Poultry Biology Reviews* 11:11-20.
159. Krushinskie EA. 2006. U.S. poultry industry preparedness for an HPAI outbreak. Avian influenza: dealing with the challenge. bulldogsolutions.net/WattPublishing/WPC01102006/frmEventDescription.aspx. Accessed April 1, 2007.
160. Cutler GJ. 1986. The nature and impact of layer industry changes. *United States Animal Health Association. Second International Symposia on Avian Influenza*, pp. 423-6.
161. Alexander DJ. 1993. Orthomyxovirus infection. In: McFerran JB and McNulty MS (eds.), *Virus Infections of Birds* (Amsterdam, The Netherlands: Elsevier Science Publishers, pp. 287-316).
162. Canning K. 2005. A matter of pride. *Refrigerated and Frozen Foods*. www.refrigeratedfrozenfood.com/content.php?s=RF/2005/12&p=8. Accessed April 1, 2007.
163. Schmit J. 2005. Poultry farm tactics may thwart bird flu. *USA Today*, November 14. usatoday.com/news/nation/2005-11-13-farmers-birdflu_x.htm?csp=N009. Accessed April 1, 2007.
164. Tablante NL, San Myint M, Johnson YJ, Rhodes K, Colby M, and Hohenhaus G. 2002. A survey of biosecurity practices as risk factors affecting broiler performance on the Delmarva Peninsula. *Avian Diseases* 46:730-4.
165. Beard CW. 2003. Minimizing the vulnerability of poultry production chains for avian influenza. In: Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control (Wageningen, The Netherlands, pp. 133-7).
166. Beard CW. 2003. Minimizing the vulnerability of poultry production chains for avian influenza. In: Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control (Wageningen, The Netherlands, pp. 133-7).
167. Corporate Crime Reporter. 2003. Dirty Money: Corporate Criminal Donations to the Two Major Parties. July 3. www.corporatecrimereporter.com/ccrreport.pdf. Accessed April 1, 2007.
168. Tyson Foods. 2005. Tyson provides \$26 million to fuel family farmers: supplemental energy allowance addresses rising fuel costs. Tyson Press Room. October 3. www.tyson.com/Corporate/PressRoom/ViewArticle.aspx?id=1925. Accessed April 7, 2007.
169. 2005. *IBP, Inc. v. Alvarez* (03-1238); *Tum v. Barber Foods, Inc* (04-66) Supreme Court collection. www.law.cornell.edu/supct/cert/03-1238.html. Accessed April 1, 2007.
170. Schmit J. 2005. Poultry farm tactics may thwart bird flu. *USA Today*, November 14. usatoday.com/news/nation/2005-11-13-farmers-birdflu_x.htm?csp=N009. Accessed April 1, 2007.
171. Animal Health and Welfare Panel of the European Food Safety Authority. 2005. Animal health and welfare aspects of avian influenza. *European Food Safety Authority Journal* 266:1-21.
172. Beard CW. 2003. Minimizing the vulnerability of poultry production chains for avian influenza. In: Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control (Wageningen, The Netherlands, pp. 133-7).
173. Stegeman A. 2003. Workshop 1: introduction and spread of avian influenza. In: Schrijver RS and Koch G (eds.), Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control. library.wur.nl/frontis/avian_influenza/workshop1.pdf. Accessed April 1, 2007.
174. Shane SM. 2003. Disease continues to impact the world's poultry industries. *World Poultry* 19(7):22-7.
175. Rudd K. 1995. Poultry reality check needed. *Poultry Digest*, December, pp. 12-20.
176. Rudd K. 1995. Poultry reality check needed. *Poultry Digest*, December, pp. 12-20.

177. Anderson I. 2002. Foot and mouth disease 2001: lessons to be learned inquiry report (London, UK: The Stationery Office).
178. Shane SM. 2003. Disease continues to impact the world's poultry industries. *World Poultry* 19(7):22-7.
179. Plumiers F. 2003. Workshop 4: control measures and legislation. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control* (Wageningen, The Netherlands). library.wur.nl/frontis/avian_influenza/workshop4.pdf. Accessed April 1, 2007.
180. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
181. Thaxton YV. 2005. Are you prepared for AI? *Poultry*, April/May, p. 5.
182. Centers for Disease Control and Prevention. 2004. Update on avian influenza A (H5N1). April 12. www.cdc.gov/flu/avian/professional/han081304.htm. Accessed April 1, 2007.
183. Richmond JY. 1998. The 1, 2, 3's of Biosafety Levels. Centers for Disease Control and Prevention Office of Health and Safety, February 6. www.cdc.gov/od/ohs/symp5/jyrtext.htm. Accessed April 1, 2007.
184. Centers for Disease Control and Prevention. 2004. Update on avian influenza A (H5N1). April 12. www.cdc.gov/flu/avian/professional/han081304.htm. Accessed April 1, 2007.
185. Environment, Health and Safety Division: Ernest Orlando Lawrence Berkeley National Laboratory. Biosafety program.
186. Richmond JY. 1998. The 1, 2, 3's of Biosafety Levels. Centers for Disease Control and Prevention Office of Health and Safety, February 6. www.cdc.gov/od/ohs/symp5/jyrtext.htm. Accessed April 1, 2007.
187. Environment, Health and Safety Division: Ernest Orlando Lawrence Berkeley National Laboratory. Biosafety program.
188. Richmond JY. 1998. The 1, 2, 3's of Biosafety Levels. Centers for Disease Control and Prevention Office of Health and Safety, February 6. www.cdc.gov/od/ohs/symp5/jyrtext.htm. Accessed April 1, 2007.
189. Environment, Health and Safety Division: Ernest Orlando Lawrence Berkeley National Laboratory. Biosafety program.
190. Stegeman A. 2003. Workshop 1: introduction and spread of avian influenza. In: Schrijver RS and Koch G (eds.), *Proceedings of the Frontis Workshop on Avian Influenza: Prevention and Control*. library.wur.nl/frontis/avian_influenza/workshop1.pdf. Accessed April 1, 2007.
191. 2006. Live birds back on sale despite ban. *Shanghai Daily News*, March 2. english.eastday.com/eastday/englishedition/node20665/node20669/node22813/node95960/node95963/node95964/node95966/userobject1ai1885583.html. Accessed April 1, 2007.
192. Hoo S. 2005. China closes all Beijing poultry markets. *Associated Press*, November 7. katv.com/news/stories/1105/275446.html. Accessed April 1, 2007.
193. 2006. Hong Kong to ban live poultry sales in three years. *Agence France Presse*. news.yahoo.com/s/afp/healthfluhongkong:_ylt=Ak6dh2DxZxiPZHnhKChGjwiTvyli:_ylu=X3oDMTBiMW04NW9mBHNIYwMIJVRPUCU. Accessed April 1, 2007.
194. 2006. Beijing closes live poultry market permanently. *ThePoultrySite.com* December 18. www.thepoultrysite.com/poultrynews/10522/beijing-closes-live-poultry-market-permanently. Accessed April 1, 2007.
195. 2006. Law to ban slaughtering live poultry in markets. *Taiwan News*, August 31. english.www.gov.tw/TaiwanHeadlines/index.jsp?print=1&catid=10&recordid=98885. Accessed April 1, 2007.
196. Environmental News Service. 2005. Beijing halts poultry trade to guard against bird flu. November 7. www.ens-newswire.com/ens/nov2005/2005-11-07-05.asp. Accessed April 1, 2007.
197. Chuang J. 2006. Anti-bird flu plan will cost poultry vendors jobs. *Taipei Times*, August 31, p. 1.
198. 2006. Law to ban slaughtering live poultry in markets. *Taiwan News*, August 31. english.www.gov.tw/TaiwanHeadlines/index.jsp?print=1&catid=10&recordid=98885. Accessed April 1, 2007.
199. 2003. Chicken flu cast doubts on old eating habits in HK. *Shanghai Star*, March 27. app1.chinadaily.com.cn/star/2003/0327/fe22-2.html. Accessed April 1, 2007.
200. Woo PC, Lau SK, and KY Yuen. 2006. Infectious diseases emerging from Chinese wet-markets: zoonotic origins of severe respiratory viral infections. *Current Opinions in Infectious Disease* 19(5):401-7.

201. Webster RG and Hulse DJ. 2004. Microbial adaptation and change: avian influenza. *Revue Scientifique et Technique* 23(2):453-65.
202. Webby RJ and Webster RG. 2001. Emergence of influenza A viruses. *Philosophical Transactions of the Royal Society of London* 356:1817-28.
203. Choi YK, Seo SH, Kim JA, Webby RJ, and Webster RG. 2005. Avian influenza viruses in Korean live poultry markets and their pathogenic potential. *Virology* 332:529-37.
204. Lashley FR. 2004. Emerging infectious diseases: vulnerabilities, contributing factors and approaches. *Expert Review of Anti-Infective Therapy* 2(2):299-316.
205. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
206. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
207. U.S. Department of Agriculture Agricultural Research Service. Action plan.
208. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
209. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
210. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
211. Webster RG. 2004. Wet markets: a continuing source of severe acute respiratory syndrome and influenza? *Lancet* 363:234-6.
212. Swayne DE. 2003. Transcript of the question and answer session from the Fifth International Symposium on Avian Influenza. *Avian Diseases* 47:1219-55.
213. Suarez DL, Spackman E, and Senne DA. 2003. Update on molecular epidemiology of H1, H5, and H7 influenza virus infections in poultry in North America. *Avian Diseases* 47:888-97.
214. Mullaney R. 2003. Live-bird market closure activities in the northeastern United States. *Avian Diseases* 47:1096-8.
215. Associated Press. 2004. Bird flu found at four New Jersey live chicken markets. *USA Today*, February 12. usatoday.com/news/nation/2004-02-12-bird-flu_x.htm. Accessed April 1, 2007.
216. Kingsbury K. 2004. Animal markets are alive with controversy. *Columbia News Service*. February 16. www.jrn.columbia.edu/studentwork/cns/2004-02-16/455.asp. Accessed April 1, 2007.
217. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
218. Bulaga LL, Garber L, Senne D, et al. 2003. Descriptive and surveillance studies of suppliers to New York and New Jersey retail live-bird markets. *Avian Diseases* 47:1169-76.
219. U.S. Department of Agriculture Agriculture Research Service. 2005. Science Hall of Fame. December 16. www.ars.usda.gov/careers/hof/browse.htm. Accessed April 1, 2007.
220. Thacker C. 2006. Live markets: a risk for entry of bird flu into US. *Dow Jones Newswires*, March 31. cattlenetwork.com/content.asp?contentid=26782. Accessed April 1, 2007.
221. U.S. Department of Agriculture Food Safety Inspection Service. 2005. Guidance for determining whether a poultry slaughter or processing operation is exempt from inspection requirements of the Poultry Products Inspection Act. June.
222. Iowa Department of Agriculture and Land Stewardship Meat and Poultry Inspection Bureau. Slaughter-processing-labeling-marketing: the basics. www.agriculture.state.ia.us/thebasics.htm. Accessed April 1, 2007.

223. Shane S. 2004. Live-bird markets are under the microscope: as the United States battles new outbreaks of bird flu, the role and necessity of live-bird markets must be examined. *National Provisioner* 218(4):38. www.findarticles.com/p/articles/mi_hb314/is_200404/ai_hibm1G1117180028. Accessed April 1, 2007.
224. Personal communication with Patrice Klein, DVM, Staff Veterinarian/Avian Disease Specialist, USDA/APHIS; Fidelis Hegngi, DVM, Senior Staff Veterinarian, USDA/APHIS; Thomas J. Myers, DVM, Co-Director of the USDA's National Center for Animal Health Programs; and Ed Curlett, spokesperson for USDA, on July 26, 2006.
225. Senne DA, Suarez DL, Stallnecht E, Pedersen JC, and Panigrahy B. 2006. Ecology and epidemiology of avian influenza in North and South America. *Developments in Biologicals* 124:37-44.
226. Hedley AJ, Leung GM, Fielding R, and Lam TH. 2004. The prevention of avian influenza in Hong Kong: observations on the HKSAR government's consultation document. Department of Community Medicine and Unit for Behavioural Sciences, School of Public Health, The University of Hong Kong. legco.gov.hk/yr03-04/english/panels/fseh/papers/fe0604cb2-2583-03-e.pdf. Accessed April 1, 2007.
227. Webby RJ and Webster RG. 2001. Emergence of influenza A viruses. *Philosophical Transactions of the Royal Society of London* 356:1817-28.