

WellBeing International

WBI Studies Repository

11-27-2013

Stress and Reproductive Hormones in Grizzly Bears Reflect Nutritional Benefits and Social Consequences of a Salmon Foraging Niche

Heather M. Bryan
University of Victoria

Chris T. Darimont
University of Victoria

Paul C. Paquet
University of Calgary

Katherine E. Wynne-Edwards
University of Calgary

Judit E. G. Smits
University of Calgary

Follow this and additional works at: <https://www.wellbeingintludiesrepository.org/physio>



Part of the [Animal Studies Commons](#), [Other Animal Sciences Commons](#), and the [Veterinary Physiology Commons](#)

Recommended Citation

Bryan HM, Darimont CT, Paquet PC, Wynne-Edwards KE, Smits JEG (2013) Stress and Reproductive Hormones in Grizzly Bears Reflect Nutritional Benefits and Social Consequences of a Salmon Foraging Niche. PLoS ONE 8(11): e80537. <https://doi.org/10.1371/journal.pone.0080537>

This material is brought to you for free and open access by WellBeing International. It has been accepted for inclusion by an authorized administrator of the WBI Studies Repository. For more information, please contact wbisr-info@wellbeingintl.org.



Stress and Reproductive Hormones in Grizzly Bears Reflect Nutritional Benefits and Social Consequences of a Salmon Foraging Niche

Heather M. Bryan^{1,2,*}, Chris T. Darimont^{2,3}, Paul C. Paquet^{2,4}, Katherine E. Wynne-Edwards¹, Judit E. G. Smits¹

1 Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada, **2** Raincoast Conservation Foundation, Bella Bella, British Columbia, Canada, **3** Department of Geography, University of Victoria, Victoria, British Columbia, Canada, **4** Faculty of Environmental Design, University of Calgary, Calgary, Alberta, Canada

Abstract

Physiological indicators of social and nutritional stress can provide insight into the responses of species to changes in food availability. In coastal British Columbia, Canada, grizzly bears evolved with spawning salmon as an abundant but spatially and temporally constrained food source. Recent and dramatic declines in salmon might have negative consequences on bear health and ultimately fitness. To examine broadly the chronic endocrine effects of a salmon niche, we compared cortisol, progesterone, and testosterone levels in hair from salmon-eating bears from coastal BC ($n = 75$) with the levels in a reference population from interior BC lacking access to salmon ($n = 42$). As predicted, testosterone was higher in coastal bears of both sexes relative to interior bears, possibly reflecting higher social density on the coast mediated by salmon availability. We also investigated associations between the amount of salmon individual bears consumed (as measured by stable isotope analysis) and cortisol and testosterone in hair. Also as predicted, cortisol decreased with increasing dietary salmon and was higher after a year of low dietary salmon than after a year of high dietary salmon. These findings at two spatial scales suggest that coastal bears might experience nutritional or social stress in response to on-going salmon declines, providing novel insights into the effects of resource availability on fitness-related physiology.

Citation: Bryan HM, Darimont CT, Paquet PC, Wynne-Edwards KE, Smits JEG (2013) Stress and Reproductive Hormones in Grizzly Bears Reflect Nutritional Benefits and Social Consequences of a Salmon Foraging Niche. PLoS ONE 8(11): e80537. doi:10.1371/journal.pone.0080537

Editor: Nei Moreira, Federal University of Parana (UFPR) – Campus Palotina, Brazil

Received: June 18, 2013; **Accepted:** October 14, 2013; **Published:** November 27, 2013

Copyright: © 2013 Bryan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: For funding, the authors gratefully acknowledge support from: Raincoast Conservation Foundation (www.raincoast.org), Animal Welfare Institute (www.awionline.org), Blue Planet Links (www.blueplanetlinks.com), Explorers Club (www.explorers.org), Habitat Conservation Trust Foundation (www.hctf.ca), McLean Foundation (www.mcleanfoundation.ca), Moore Foundation (www.moore.org), Norcross Foundation (www.norcrossws.org), Spirit Bear Research Foundation (www.spiritbearfoundation.com), Tides Canada (tidescanada.org), Willow Grove Foundation (www.willowgrovefoundation.com) and the Wilburforce (www.wilburforce.org) Foundations, Environment Canada Science Horizons (www.ec.gc.ca/sci_hor), the University of Calgary (www.ucalgary.ca), and National Science and Engineering Council (NSERC) (<http://www.nserc-crsng.gc.ca>) Discovery grants to CTD (435683), KEW (RGPIN-106386-2008) and JES (RGPIN-22876-20). HMB was supported by a National Science and Engineering Council (NSERC) postgraduate fellowship as well as by the University of Calgary. CTD recognizes support from the Tula Foundation (www.tula.org). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: heabry@gmail.com

‡ Current address: Department of Geography, University of Victoria, Victoria, British Columbia, Canada

Introduction

Understanding the physiological responses of organisms to stressors is essential in predicting long-term consequences of environmental change [1–3]. Food limitation is a stressor that can directly affect population productivity by altering survival and reproduction [4]. Moreover, the distribution, abundance, and quality of food affect populations by mediating social structure and behavior [5]. Monitoring physiological indicators of nutritional and social stress may provide an early warning of population-level responses to environmental change [6]; this approach is particularly valuable in taxa such as ursids where long-term population productivity is difficult or impossible to quantify.

Hair provides an excellent approach for examining physiological responses to food resource shortages as it can be chemically analyzed to determine both diet and steroid hormone levels. In contrast with serum and feces, which are commonly used for measuring steroid hormones in wildlife and reflect time periods of

minutes to hours, hair reflects endocrine activity integrated over several months. Consequently, steroids in hair are insensitive to short-term stressors [7] and can be related to longer-term life history events and stages [8]. Steroid hormones are incorporated into growing hair via the blood vessel that feeds the hair follicle and/or from the follicle itself, which can synthesize steroids locally [8–10].

An increasing number of studies, most focusing on the glucocorticoid stress hormone, cortisol, have shown that steroid measurements from hair provide biologically meaningful information in humans, captive animals and wildlife [7,11–18]. Recently, several studies have provided biological validation in ursids, including grizzly bears [17,19], polar bears (*Ursus maritimus*) [16,18,20,21], and Asiatic black bears (*Ursus thibetanus*) [22]. Notably, MacBeth [23] found relatively high levels of cortisol in hair from an emaciated grizzly and an emaciated black bear (*Ursus americanus*) compared with 151 other grizzly bears. Similarly,

Malcolm et al. [22] documented higher cortisol in hair of Asiatic black bears kept under stressful conditions on a bile farm and those recently admitted to a shelter compared with bears already living at the shelter. Moreover, paired samples showed that cortisol in hair decreased as bears acclimatized to the shelter [22].

As a long-lived species that is acutely sensitive to large-scale anthropogenic disturbances [24,25], grizzly bears serve as a model system for understanding the physiological effects of food resource declines. For millennia, grizzly bears (*Ursus arctos*) in coastal British Columbia, Canada, and beyond have evolved with abundant Pacific salmon (*Oncorhynchus* spp.) that becomes available each year during the autumn spawning event [26,27]. Salmon allows bears to meet their energetic requirements more efficiently than a diet of plants alone [28–30]. Moreover, nutrients from salmon come at a critical time before hibernation when pre-denning fat reserves are positively correlated with over-winter survival and reproduction in the following year [31–33]. Among populations, grizzly bears with access to salmon have higher population density, body size and litter size [34].

Historically, salmon returns have been a relatively predictable annual event for coastal bears, though the number and timing of spawners varies among years and streams [35]. Despite some exceptions, there have been widespread or regional declines in salmon abundance through much of coastal British Columbia [36–38]. Today, fewer than 4% of streams monitored in coastal BC consistently meet their salmon escapement targets (i.e., number of salmon that escape human fishing nets and return to their natal streams to spawn) [39]. Notably, the hair of bears in North America grows for approximately six months from spring to fall [40–42], during which salmon are consumed for three months [Table 1].

To examine whether endocrine levels are potentially influenced by variation in salmon availability and consumption, we compared hormone levels in a population of coastal bears with access to salmon with an interior population without access to salmon. Among coastal bears only, we examined the relationship between hormones and salmon consumption (determined by stable isotope analysis). Between regions, we predicted that cortisol, as a general indicator of physiological stress, would be elevated in response to nutritional stress [43,44] or social instability [45]. Among coastal bears, we predicted a negative relationship between cortisol and salmon consumption, reflecting either a nutritional or social benefit of access to more salmon.

To date, no studies have examined testosterone and progesterone in bear hair. Testosterone plays an important role in reproduction and also varies in relation to the social competitive environment above levels required for reproduction [46–49]. In particular, testosterone facilitates behavioral and physical traits necessary to win social conflicts in fitness-enhancing situations [49,50]. Therefore, we predicted that testosterone would be

elevated in coastal bears, where population density is higher and social interactions occur over temporally and spatially constrained salmon runs. Among coastal males, we predicted higher testosterone in males that consume more salmon, possibly reflecting a nutritional benefit of eating salmon or higher social density in areas where more salmon is available to be eaten.

Progesterone, which is elevated in females during pregnancy and pseudopregnancy, should be positively associated with population-level reproductive activity [51], because hair grows over the time interval that incorporates follicular development, ovulation, and mating. Given the higher productivity of bear populations with access to salmon [29], we predicted that progesterone would be higher in coastal compared with interior females.

Materials and Methods

Ethics statement

Samples were collected under animal care protocols approved by the Chancellor’s Animal Research Committee at the University of California Santa Cruz (WILMc0904) and the Animal Care Committee at the University of Calgary (B110R-01). Our sampling sites occurred in the traditional territory of the Heiltsuk Nation as well as in provincial parks. Permission to collect samples from these areas was granted by the Heiltsuk Integrated Resource Management Department and BC Parks (Park Use Permit Number 103586).

Study areas and sample collection

We collected bear hair samples from coastal and interior BC (Fig. 1). On the coast, our core study area was located near Bella Bella (52°13’15.8”N, 127°45’28.4”W) where we collected hair samples using standard, grid-based DNA mark-recapture methods [52–54]. In a 2009 pilot year, we sampled over 2500 km² at 92 barbed-wire hair-snagging stations placed within 5×5 km grid cells. In 2010 and 2011, we expanded the area to 5000 km² with 71 snag stations in 7×7 km cells. We obtained additional hair from archived samples of grizzly bears killed in coastal BC in the springs of 2004–2009. These samples came from a larger area extending from Knight Inlet in the south (50°29’44.5”N, 131°36’30.5”W) to the Khutzeymateen (54°59’28.6”N, 122°36’13.8”W) grizzly bear management unit in the north [55]. Coastal bears assimilate a substantial portion of their yearly dietary protein from salmon [56]. These bears inhabit the coastal western hemlock biogeoclimatic zone of BC, which is characterized by high precipitation (average 2228 mm/year) and a temperature averaging 8°C [57].

For comparison with coastal bears, we obtained archived hair from bears in the interior of BC. The sampling extent ranged from the Moberly grizzly bear management unit in the south

Table 1. Approximate time line of hair growth and corresponding yearly natural history events of grizzly bears.

	May-October	November-April
Hair Growth	Hair grows, incorporating steroid hormones & isotopic dietary information [40–42]. May: old hair shed, new coat starts to grow. ¹	Hair stops growing over winter & does not incorporate steroid hormones or dietary information.
Bear Biology	May-July: Bears eat mainly vegetation & some terrestrial meat [61,105]. Breeding season, increases in male-male aggressive interactions[79]. Aug-Oct: Coastal bears aggregate on salmon streams. Bears gain body mass for over-winter survival & reproduction [31–33].	Hibernation. Cubs born. Bears may lose >30% of their body mass [31]

¹All hair samples collected in May were grown in the previous year.
doi:10.1371/journal.pone.0080537.t001




Figure 1. Bear hair collection areas. We collected grizzly bear hair samples (circles) from coastal and interior British Columbia (BC) in the springs of 2004–2011. The samples came either from government archives of hunted bears or hair snagging stations on the central coast of BC. The bottom left inset shows the coastal study area where we sampled from hair snagging stations in springs of 2009–2011. doi:10.1371/journal.pone.0080537.g001

(55°32'25.4"N, 129°37'30.6"W) to the Hyland and Muskwa units in the north (59°46'8.6"N, 126°40'2.3"W). Bears in these regions eat plants and terrestrial meat and do not have access to anadromous salmon or kokanee [56]. In contrast with coastal bears, interior bears inhabit a region with lower precipitation (330–570 mm/year) and a continental climate characterized by warm summers, cool winters, and an average yearly temperature between -3 and 3°C [57,58].

Regarding human interactions, both bear populations have coexisted with local First Nations for thousands of years; today, the human population density in both areas is relatively low compared with elsewhere in the province [58,59]. Industrial activities such as logging occur in both areas but the extent of human activity, particularly road density, is higher in the interior [57,60].

All hair samples were collected in spring and therefore reflected hair grown from spring to fall of the previous year (Table 1). Notably, bear hair grows—and incorporates hormones—at approximately one cm/month over the six month period when bears are most active [61]. We stored the samples in paper envelopes at room temperature in a dark, dry environment [28].

Genetic Analyses

After collection, we sent all samples to a commercial laboratory (Wildlife Genetics International, Nelson, BC, Canada) where seven microsatellite markers were used to identify individual bears as well as their species and sex [62]. We used the remaining hair shafts for stable isotope analysis and hormone assays. When multiple samples collected from single or different snag stations

were identified genetically as being from one individual in the same year, we pooled samples to obtain enough material for hormone assays.

Stable isotope analysis to quantify salmon consumption

We prepared samples for stable isotope analysis as previously described [63,64]. Subsequently, we sent the samples to the University of Saskatchewan’s stable isotope facility where the ratios of nitrogen (¹⁵N/¹⁴N) and carbon (¹³C/¹²C) stable isotopes were measured using gas chromatography mass spectrometry. To estimate the proportion of salmon in the diet of each bear, we used a Bayesian mixing model [65,66]. Following other studies of coastal grizzly bears, we assumed that bears’ diets consist only of plant or salmon-based protein [56]. We used previously published estimates of anadromous salmon and plant stable isotope signatures, standard deviations, and fractionation rates [56].

Analysis of steroids in hair

Our protocol for analyzing steroids in hair was similar to that previously published [67]. Additional details on hormonal assays and validations are provided as supporting information (Fig. S1, Table S1, Text S1).

Bear density estimates

Across the province, we classified bear density in coastal and interior bears based on government estimates for each of the grizzly bear management units of BC [55]. In the grid-based coastal study area, we divided the study area into 10 units of 342–900 km² each

based on BC's conservation landscape units [68]. We estimated bear density within a landscape unit by dividing the number of genetically unique individuals detected in 2010 and 2011 by the number of hair snag stations (all sampled with the same effort) in that landscape unit. We used this approach to provide a generalized indicator of density based on bear detections at several nearby snag stations. Therefore, we used number of hair snags rather than unit area as our denominator in this calculation. Bear density estimates fit naturally into high and low classifications.

Data Analysis

All analyses were conducted in R [69]. Before analysis, we removed outliers falling >2 SD from the mean for cortisol and/or testosterone ($n=4$). Three of these samples had extreme values that were 15–37 times higher than the mean and were beyond the range of the hormone assays prior to dilution. The fourth sample was a multivariate outlier for cortisol and testosterone, identified using the gap test [70]. In future, and if similar outliers are a consistent finding, they may be an interesting subset to consider, possibly reflecting high physiological stress. However, they might also be caused by extreme cases of external contamination not removed by our wash procedure or an unidentified error in the laboratory. Here, we assumed the latter possibility and excluded these individuals from statistical analysis. One coastal bear was excluded for having a non-coastal dietary salmon signature; we suspected this was due to an error or mix-up in the database. To improve normality, we applied a negative reciprocal transformation to cortisol and testosterone and an arcsin transformation to our proportion of salmon in diet metric [71]. Progesterone did not require a transformation, possibly because of few samples from female bears ($n=21$). We used t-tests to check for differences in samples collected from hunters and hair snags in coastal BC.

Comparison of coastal and interior bear populations. We developed candidate linear regression models to examine the effects of region, sex, and bear density on cortisol and testosterone. Model sets for both hormones included an interaction between region and sex because we predicted that males and females might respond differently to salmon availability across regions as well as a null model containing a constant. We ranked models using Akaike information criterion, corrected for small sample size (AICc), and considered our top model set to include all candidate models with a Δ AICc score <2 . To assess the adequacy of top models, we plotted histograms of the residuals, residuals versus predictors and residuals versus predicted values. In addition, we examined Cook's distance as an indicator of influential observations. Variance inflation factors for top models ranged from 1.0 to 2.0 indicating low collinearity among variables. For comparisons of progesterone, we used a linear model to compare coastal ($n=15$) to interior females ($n=9$) and a t-test to examine differences between females ($n=21$) and males ($n=4$).

Salmon-hormone relationships in coastal male bears. For these analyses, we excluded female bears because of their small sample size ($n=15$ of 70 bears). We first explored our prediction of a direct relationship between cortisol or testosterone and salmon consumption among all coastal bears ($n=55$) using linear regression. In addition, we posited that the amount of salmon bears consumed in autumn would influence their nutritional and physiological state emerging from hibernation in the following year; therefore, we examined whether there was a lag between hormone levels and salmon consumption by examining trends over time. To address this question, we focused on cortisol and testosterone in coastal males from our grid-based study area ($n=28$; where we had a field-based measure of relative density) to compare the effects of salmon consumption, year, and

bear density using linear models, as described above. We centred and scaled the salmon consumption metric so that parameter estimates of variables would be comparable. Variance inflation factors for all models ranged from 1.0 to 1.4. Finally, we used f-tests and paired t-tests to examine trends in individuals detected (and measured with hormonal and isotopic assays) in two years of the study ($n=7$).

Results

General patterns

Overall, the median cortisol concentration was 8.1 pg/mg [range: 5.3–26.1] in 113 hair samples, the median testosterone was 5.6 pg/mg [range: 3.1–21.1] in 112 samples, and the median progesterone was 26.2 pg/mg [range: 9.1–46.2] in 27 samples. Among coastal bears, cortisol and testosterone were similar in hair samples collected from hunters and snag stations so samples were pooled in subsequent analyses (cortisol: $t=0.27$, $df=66$, $p=0.79$; testosterone: $t=-0.23$, $df=67$, $p=0.82$).

Comparison of hormones in coastal and interior bear populations

The most striking difference between regions was higher testosterone in coastal bears of both sexes, which is consistent with our prediction of higher social density among bears with access to salmon (Fig. 2B; Table 2). As expected, the top model also revealed higher testosterone in males than females (Fig. 2B; Table 2). By contrast, the top model for cortisol was the null model, suggesting no differences between coastal and interior bear populations or sexes (Fig. 2A; Table 2). Notably, bear density was not an important predictor of cortisol or testosterone (Table 2). Progesterone did not differ between regions (Fig. 2C; Table 2) and was higher in females ($t=-6.2$, $df=15$, $p<0.001$; Fig. 2C).

Salmon-hormone relationships in coastal male bears

As predicted among all coastal males ($n=55$), hair cortisol decreased with increasing dietary salmon, though very marginally (adj $R^2=0.06$, $F_{1,53}=4.2$, $p=0.046$; Fig. 3A). In the smaller coastal study area that we sampled consistently in three years ($n=28$), our model selection approach identified the effects of salmon consumption and year as being important predictors of cortisol (Table 3). Similar to the trend in the larger dataset, cortisol decreased marginally with increasing salmon consumption. Moreover, the top model set revealed that cortisol was higher in 2008 and 2009 after years of low average salmon consumption compared with 2010 after a year of higher salmon consumption (Fig. 3B; Table 3).

In contrast with our prediction, there was no evidence of a relationship between testosterone and dietary salmon among all coastal males (adj $R^2=0$, $F_{1,53}=0.97$, $p=0.33$). In the grid-based coastal study area, the top model for testosterone was the null, revealing that the variables we examined explained little of the variability in hair testosterone. However, the top model set included weak effects of bear density and salmon consumption (Table 3). In contrast with our predictions, testosterone decreased with increasing salmon consumption and was lower in areas of high bear density.

Trends in the seven bears sampled in both years reflected those at the population level; these bears had more variable cortisol in 2009 than in 2010 ($F_{5,5}=25.3$, $p=0.003$; Fig. 4A). Cortisol levels in several bears were lower in 2010 than 2009, but the difference was not significant (paired $t=1.80$, $df=5$, $p=0.13$). Testosterone did not show a consistent trend between years ($t=0.03$, $df=5$, $p=0.98$; Fig. 4B).




Figure 2. Mean and standard error for cortisol (A), testosterone (B), and progesterone (C) across regions and sexes. (A) Cortisol was similar between sexes and regions in grizzly bears. (B) Testosterone was higher in males and coastal bears. (C) Progesterone was higher in females. Cortisol and testosterone were reciprocal transformed (-1/x) to improve normality; progesterone is expressed in pg/mg of hair. Sample sizes are displayed below the error bars. doi:10.1371/journal.pone.0080537.g002

Discussion

General patterns

Hormone measurements in hair—which reflect long-term endocrine information—provide novel insights into the physiological responses of wildlife to environmental change [8,16–18,20,21]. Interestingly, the cortisol values in our study (median:

8.1 pg/mg, range: 5.3–26.1) were higher than those previously reported in 151 live-captured grizzly bears from Alberta, Canada (median: 2.8 pg/mg, range: 0.6–43.3 pg/mg; [17]). These differences are probably methodological but could also relate to differences in population densities, habitat, genetics, or the dynamics of natural and anthropogenic stressors.

Our findings show that immunoreactive progesterone and testosterone can be measured relatively easily in addition to cortisol from the same hair sample. Moreover, hormone concentrations revealed expected differences between sexes. Elevated testosterone in hair of males reflects higher testosterone levels in males during the breeding season, which occurs in June and July [72]. Similarly, progesterone is elevated in female bears following fertilization, which occurs in spring or early summer [73]. Though the corpora lutea are mostly dormant until implantation occurs in late fall, they produce enough progesterone that levels are elevated above baseline in pregnant and pseudo-pregnant females [74,75].

Comparison of coastal and interior bear populations

This study also provides insight into the physiological implications of living in an environment with a nutritious but seasonally and spatially constrained resource. Salmon provides nutritional benefits to bears [31,32,34]; however, higher testosterone in coastal bears could have fitness costs such as increased energetic expenditure and risk of injury from intraspecific interactions, as well as impaired immunity [47,48]. Differences in testosterone between coastal and interior bears could also relate directly to diet. Spawning salmon have high levels of androgens, which could potentially increase circulating testosterone levels in bears, which have access to salmon for approximately three of the six months of hair growth [76]. However, we found no support for this possibility as dietary salmon and testosterone were not correlated in coastal males. Moreover, cortisol was not higher in coastal bears, even though spawning salmon have extremely high levels of glucocorticoids [77].

The higher testosterone levels of coastal bears might relate to their larger body size compared with interior bears. Previous studies of bears have shown that testosterone is positively linked with body size in males during the breeding season [78,79]; however, it is not clear whether the same trend would occur between populations differing in average size or whether the relationship would be detectable in hair, which integrates endocrine activity during the breeding and non-breeding periods.

Our findings show that testosterone did not vary with provincial estimates of bear density. It is possible that the spatiotemporal distribution of resources at scales smaller than region is a more important mediator of social interactions than the number of bears in a region [80,81]. Indeed, several characteristics of habitat and resource availability would affect the frequency and type of social interactions (i.e., social density) between regions. Whereas interior bears use a variety of habitats from the treeline to the alpine [58,82], coastal bears spend most of their time along valley bottoms due to less usable habitat on the coast [83]. Food sources also differ; interior bears feed on vegetation and opportunistically on ungulates [58,80,84]. These bears use productive habitats such as burns and berry patches but the feeding aggregations are less pronounced compared with those on spawning salmon streams [85,86]. Indeed, well-described social interactions over access to salmon often lead to aggressive encounters and the establishment of dominance hierarchies [80,86]. An influx of bears from the interior to salmon spawning streams in the fall would make the social dynamics particularly intense.

In coastal female bears, higher testosterone compared with interior females might reflect higher reproductive rates. However,

Table 2. Estimates and standard error (in parentheses) of parameters in the top models ($\Delta AIC_c \leq 2$) describing cortisol, testosterone and progesterone in hair of coastal ($n = 70$) and interior ($n = 42$) grizzly bears.

Response (models compared)	Intercept	Sex = Female	Region = Interior	Sex* Region	Bear Density ^a
Cortisol (10)	-0.119* (0.003)	-	-	-	-
Testosterone (8)	-0.161* (0.007)	-0.031* (0.013)	-0.028* (0.011)	-	-
Progesterone ^b (2)	29.2* (1.67)	5 ^c	-	NE ^d	NE ^d

*Significant at $\alpha = 0.05$

^aBased on government estimates of density in grizzly bear management units

^bFemales only

^cSignificant, a separate t-test showed that females ($n = 21$) had higher progesterone than males ($n = 4$)

^dNot examined

doi:10.1371/journal.pone.0080537.t002

this is unlikely the only explanation as the latter stages of pregnancy—when testosterone levels are highest—occur in winter when hair is not growing and does not incorporate steroid hormones. Alternatively, higher testosterone in coastal females might be modulated by social conditions. Though usually considered with respect to reproductive traits in males, testosterone in females has been linked with defending resources and acquiring food [87,88]. In female bears, elevated testosterone might be advantageous in obtaining salmon, a food that increases reproductive success but that can be difficult to obtain because of intense competition with other bears [34,86]. Testosterone in female bears could also relate to aggressive encounters to prevent infanticide by males or other females on salmon spawning streams [89,90]. Additional studies of female bears with and without cubs

would be helpful in understanding factors affecting testosterone levels in females.

In contrast with testosterone, cortisol was similar between regions, suggesting that coastal and interior bears experience similar levels of physiological stress or have different baseline cortisol levels. One possible explanation for the lack of a difference is that the nutritional benefit of access to salmon is overwhelmed by costs imposed by higher social density among coastal bears. Similarly, there was no evidence that progesterone differed between populations, possibly because we were not able to account for age, reproductive history (e.g., inter-birth interval and presence of cubs) or reproductive success (i.e., successful versus pseudo-pregnancies).




Figure 3. Evidence suggesting a relationship between cortisol and dietary salmon in male coastal grizzly bears. (A) Across all coastal males ($n = 55$), cortisol was weakly but negatively correlated with dietary salmon. (B) In the smaller, grid-based study area, mean cortisol was lower in 2010 following a year of high population-level dietary salmon than in 2009 following a year when bears ate less salmon. Note that we have no dietary salmon data from 2007, which might influence cortisol in 2008. (B). Letters above the error bars show significantly different groups. Error bars represent standard error. Sample sizes are presented below the error bars. To improve normality of residuals, cortisol was reciprocal-transformed ($-1/x$) and dietary salmon was arcsin transformed. doi:10.1371/journal.pone.0080537.g003

Table 3. Estimates and standard error (in parentheses) of parameters in the top models ($\Delta AICc \leq 2$) describing relationships between hormones and variables relating to salmon consumption among coastal male bears (n = 28).

Response (models compared)	$\Delta AICc$	Intercept	Salmon Consumption	Year ^a (2008)	Year ^a (2009)	Bear Density ^b (= High)
Cortisol (8)	0	-0.131 (0.007)	-0.011 (0.006)	0.026 (0.015)	0.030* (0.010)	-
	0.86	-0.133 (0.007)	-	0.042* (0.013)	0.027* (0.013)	-
Testosterone (8)	0	-0.161 (0.011)	-	-	-	-
	0.76	-0.125 (0.023)	-0.022 (0.012)	-	-	-0.049 (0.028)
	1.25	-0.161 (0.011)	-0.012 (0.011)	-	-	-
	1.44	-0.142 (0.022)	-	-	-	-0.025 (0.025)

*Significant at $\alpha = 0.05$

^a2010 is the reference category

^bBased on field estimates of bear density; low bear density is the reference category.

doi:10.1371/journal.pone.0080537.t003

Relationships between hormones and salmon consumption among coastal bears

Two lines of evidence support our prediction that cortisol would be higher in coastal bears that eat less salmon. Among coastal males, there was a weak but significant negative relationship between cortisol and dietary salmon. Elevated cortisol could be an adaptive response to food shortage to mobilize fat [73]. Cortisol might also play a role in bone resorption during periods of nutritional stress [91] or affect the amount and type of foods consumed [92]. Moderately elevated glucocorticoid levels could also improve foraging efficiency during reduced food availability by enhancing spatial memory [93], increasing exploratory behavior [94], or promoting innovation of novel foraging approaches [95].

The negative association between diet and cortisol could also reflect lower social tension when there are more salmon to eat. Grizzly bears in coastal areas have a social hierarchy with larger, older males being dominant over smaller, younger males [86].

Bears would be more tolerant of each other and would not have to vie for access to salmon when they are abundant [96].

Additional evidence of a direct relationship between cortisol and dietary salmon comes from our grid-based study area where we sampled consistently over three years. Cortisol was higher in 2008 and 2009 after years of low dietary salmon than in 2010 after a year of higher dietary salmon. This suggests that the amount of salmon bears consume in fall influences circulating cortisol and therefore deposition in hair in the following spring. Previous studies have established that bears entering hibernation in poor body condition have lower body mass and reproductive success in the following year [31–33]. Elevated cortisol in spring might play a role in minimizing further weight loss after a year of low dietary salmon by maximizing energy intake from low-fat, herbaceous foods, which are available in spring [97,98].

More data over several years, especially on individuals sampled multiple times, would help determine whether cortisol levels in hair relate to dietary salmon in the year of hair growth or during the spawning salmon season in the previous year. It would also be possible to segment hair corresponding to spring and fall periods in




Figure 4. Hormones in male bears detected in 2009 and 2010 reflected trends at the population level. Cortisol was generally higher and more variable in 2009 following a year of low salmon abundance compared with 2010 after a year of relatively high salmon abundance (A). Testosterone did not show a consistent trend between years (B). Each line type and point symbol represents an individual bear. No data were included from 2008 because none of the samples from recaptured bears in that year had sufficient material for hormonal analysis. doi:10.1371/journal.pone.0080537.g004

order to partition whether cortisol relates to the previous or the same-year dietary salmon [99,100]. In future studies, it will be important to monitor factors such as temperature, productivity of herbaceous foods, and precipitation, which could affect hormone levels and vary among years [101]. More studies are required to determine whether elevated cortisol has negative fitness consequences for bears, as has been shown for corticosteroids deposited into hair of polar bears [16] and feathers of sparrows [102].

Contrary to our prediction, we found only weak evidence of a relationship between testosterone and salmon consumption and the trend was opposite to our initial prediction. The negative association between testosterone and salmon consumption might occur if there is less competition for salmon when more salmon is available. This possibility could be further explored by examining the effect of salmon availability, which would influence social conditions, in addition to salmon consumption.

Conclusion

This work shows that variation in salmon abundance and consumption affects bears by altering nutritional and/or socially-mediated physiology. If salmon returns consistently decline in the future, grizzly bears that do not obtain enough salmon might experience chronically elevated cortisol and testosterone via increased nutritional and/or social stress, with unknown, but probably adverse, fitness costs. Moreover, our findings underscore the importance of considering implications for wildlife that share resources with humans as part of ecosystem-based fisheries management strategies [103,104]. Ultimately, this work adds to a growing understanding of the value of measuring stress and reproductive measures in wildlife hair as indicators of broader population health and processes [8,16,19].

Supporting Information

Figure S1 Parallelism of assay standards and diluted hair extracts for (A) cortisol, (B) testosterone, and (C)

References

- Wikelski M, Cooke SJ (2006) Conservation physiology. *Trends Ecol Evol* 21: 38–46.
- Hofer H, East ML (2012) Stress and immunosuppression as factors in the decline and extinction of wildlife populations: concepts, evidence and challenges. In: Aguirre AA, Ostfeld RS, Daszak P, editors. *New directions in conservation medicine: applied cases of ecological health*. New York, U.S.A.: Oxford University Press. pp. 109–133.
- Seebacher F, Franklin CE (2012) Determining environmental causes of biological effects: the need for a mechanistic physiological dimension in conservation biology. *Philos Trans R Soc Lond B Biol Sci* 367: 1607–1614.
- Kitaysky A, Piatt J, Wingfield J (2007) Stress hormones link food availability and population processes in seabirds. *Mar Ecol Prog Ser* 352: 245.
- Wittig RM, Boesch C (2003) Food competition and linear dominance hierarchy among female chimpanzees of the Tai National Park. *Int J Primatol* 24: 847–867.
- Buck CL, O'Reilly KM, Kildaw SD (2007) Interannual variability of Black-legged Kittiwake productivity is reflected in baseline plasma corticosterone. *Gen Comp Endocrinol* 150: 430–436.
- Ashley NT, Barboza PS, Macbeth BJ, Janz DM, Cattet MRL, et al. (2011) Glucocorticosteroid concentrations in feces and hair of captive caribou and reindeer following adrenocorticotropic hormone challenge. *Gen Comp Endocrinol* 172: 382–391.
- Meyer JS, Novak MA (2012) Hair cortisol: a novel biomarker of hypothalamic-pituitary-adrenocortical activity. *Endocrinology* 153: 4120–4127.
- Pragst F, Balikova MA (2006) State of the art in hair analysis for detection of drug and alcohol abuse. *Clin Chim Acta* 370: 17–49.
- Keckeis K, Lepschy M, Schöpfer H, Moser L, Troxler J, et al. (2012) Hair cortisol: a parameter of chronic stress? Insights from a radiometabolism study in guinea pigs. *Journal of Comparative Physiology B* 182: 985–996.
- Koren L, Mokady O, Geffen E (2006) Elevated testosterone levels and social ranks in female rock hyrax. *Horm Behav* 49: 470–477.
- Koren L, Mokady O, Karaskov T, Klein J, Koren G, et al. (2002) A novel method using hair for determining hormonal levels in wildlife. *Anim Behav* 63: 403–406.
- Martin JGA, Reale D (2008) Animal temperament and human disturbance: Implications for the response of wildlife to tourism. *Behav Process* 77: 66–72.
- Koren L, Geffen E (2009) Androgens and social status in female rock hyraxes. *Anim Behav* 77: 233–238.
- Lovari S, Pellizzi B, Boesi R, Fusani L (2009) Mating dominance amongst male Himalayan tahr: Blonds do better. *Behav Process* 81: 20–25.
- Macbeth BJ, Cattet MRL, Obbard ME, Middel K, Janz DM (2012) Evaluation of hair cortisol concentration as a biomarker of long-term stress in free-ranging polar bears. *Wildl Soc Bull* 36: 747–758.
- Macbeth BJ, Cattet MRL, Stenhouse GB, Gibeau ML, Janz DM (2010) Hair cortisol concentration as a noninvasive measure of long-term stress in free-ranging grizzly bears (*Ursus arctos*): considerations with implications for other wildlife. *Can J Zool* 88: 935–949.
- Bechshoft TØ, Sonne C, Dietz R, Born EW, Novak MA, et al. (2011) Cortisol levels in hair of East Greenland polar bears. *Sci Total Environ* 409: 831–834.
- Bourbonnais M, Nelson T, Cattet M, Darimont C, Stenhouse G (2013 accepted) Spatial analysis of factors influencing long-term stress in the grizzly bear (*Ursus arctos*) population of Alberta, Canada. *PLoS One*.
- Bechshoft TØ, Rigét FF, Sonne C, Letcher RJ, Muir DCG, et al. (2012) Measuring environmental stress in East Greenland polar bears, 1892–1927 and 1988–2009: What does hair cortisol tell us? *Environ Int* 45: 15–21.
- Bechshoft TØ, Sonne C, Dietz R, Born E, Muir D, et al. (2012) Associations between complex OHC mixtures and thyroid and cortisol hormone levels in East Greenland polar bears. *Environ Res* 116: 26–35.
- Malcolm K, McShea W, Van Deelen T, Bacon H, Liu F, et al. (2013) Analyses of fecal and hair glucocorticoids to evaluate short-and long-term stress and recovery of Asiatic black bears (*Ursus thibetanus*) removed from bile farms in China. *Gen Comp Endocrinol* 185: 97–106.
- Macbeth B (2013) An evaluation of hair cortisol concentration as a potential biomarker of long-term stress in free-ranging grizzly bears (*Ursus arctos*), polar

progesterone. Hair concentrations were measured in pg/mg of hair; regression lines were shifted on the x-axis for better visualization.

(TIF)

Table S1 Validations for analysis of bear hair using commercial cortisol, testosterone and progesterone enzyme immunoassays (Salimetrics, Philadelphia, Pennsylvania, USA).

(DOC)

Text S1 Supplementary info

(DOC)

Acknowledgments

We thank the Heiltsuk Integrated Management Department for allowing sample collection in Heiltsuk Territory. For assistance with sample collection, we are grateful to Kyle Artelle, Doug Brown, Harvey Brown, Cody Caruso, Howard Humchitt, Rosemary Invik, Ian Jansma, Collin Reid, and Christina Service. We thank Don Arney for providing helicopter support. We are grateful to William Housty for sharing samples from coastal BC. Tony Hamilton shared hair samples from BC government archives and provided input and discussion. We extend our gratitude to Lea Bond, Lee Koren, and Rosemary Invik for helping with hormonal analysis and to Monique Arseneau, Kyle Artelle, Hannah Kobluk, Steve Lever and Christina Service, for preparing isotope samples. Finally, we thank an anonymous reviewer for helpful comments.

Author Contributions

Conceived and designed the experiments: HMB CTD PCP JEGS. Performed the experiments: HMB CTD KEWE JEGS. Analyzed the data: HMB CTD JEGS KEWE. Contributed reagents/materials/analysis tools: CTD PCP KEWE JEGS. Wrote the paper: HMB CTD PCP KEWE JEGS.

- bears (*Ursus maritimus*), and caribou (*Rangifer tarandus* sp.). PhD Thesis. Saskatoon, Saskatchewan, Canada, Department of Veterinary Biomedical Sciences, University of Saskatchewan. 298 p.
24. Carroll C, Noss RF, Paquet PC, Schumaker NH (2004) Extinction debt of protected areas in developing landscapes. *Conserv Biol* 18: 1110–1120.
 25. Mattson DJ, Merrill T (2002) Extirpations of grizzly bears in the contiguous United States, 1850–2000. *Conserv Biol* 16: 1123–1136.
 26. Willson MF, Halupka KC (1995) Anadromous fish as keystone species in vertebrate communities. *Conserv Biol* 9: 489–497.
 27. Schindler DE, Scheuerell MD, Moore JW, Gende SM, Francis TB, et al. (2003) Pacific salmon and the ecology of coastal ecosystems. *Front Ecol Environ* 1: 31–37.
 28. Welch CA, Keay J, Kendall KC, Robbins CT (1997) Constraints on frugivory by bears. *Ecology* 78: 1105–1119.
 29. Hilderbrand GV, Jenkins S, Schwartz C, Hanley T, Robbins C (1999) Effect of seasonal differences in dietary meat intake on changes in body mass and composition in wild and captive brown bears. *Can J Zool* 77: 1623–1630.
 30. Rode KD, Robbins CT, Shipley LA (2001) Constraints on herbivory by grizzly bears. *Oecologia* 128: 62–71.
 31. Hilderbrand GV, Schwartz CC, Robbins CT, Hanley TA (2000) Effect of hibernation and reproductive status on body mass and condition of coastal brown bears. *J Wildl Manag* 64: 178–183.
 32. Belant JL, Kiehlend K, Follmann EH, Adam LG (2006) Interspecific resource partitioning in sympatric ursids. *Ecol Appl* 16: 2333–2343.
 33. Zedrosser A, Bellemain E, Taberlet P, Swenson JE (2007) Genetic estimates of annual reproductive success in male brown bears: the effects of body size, age, internal relatedness and population density. *J Anim Ecol* 76: 368–375.
 34. Hilderbrand GV, Schwartz CC, Robbins CT, Jacoby ME, Hanley TA, et al. (1999) The importance of meat, particularly salmon, to body size, population productivity, and conservation of North American brown bears. *Can J Zool* 77: 132–138.
 35. Quinn T, Gende S, Ruggerone G, Rogers D (2003) Density-dependent predation by brown bears (*Ursus arctos*) on sockeye salmon (*Oncorhynchus nerka*). *Can J Fish Aquat Sci* 60: 553–562.
 36. Slaney TL, Hyatt KD, Northcote TG, Fielden RJ (1996) Status of anadromous salmon and trout in British Columbia and Yukon. *Fisheries* 21: 20–35.
 37. Noakes DJ, Beamish RJ, Kent ML (2000) On the decline of Pacific salmon and speculative links to salmon farming in British Columbia. *Aquaculture* 183: 363–386.
 38. Peterman RM, Dorner B, Rosenfeld JS (2012) A widespread decrease in productivity of sockeye salmon (*Oncorhynchus nerka*) populations in western North America. *Can J Fish Aquat Sci* 69: 1255–1260.
 39. Price MHH, Darimont CT, Temple NF, MacDuffee SM (2008) Ghost runs: management and status assessment of Pacific salmon (*Oncorhynchus* spp.) returning to British Columbia's central and north coasts. *Can J Fish Aquat Sci* 65: 2712–2718.
 40. Hilderbrand GV, Farley S, Robbins C, Hanley T, Titus K, et al. (1996) Use of stable isotopes to determine diets of living and extinct bears. *Can J Zool* 74: 2080–2088.
 41. Schwartz CC, Miller SD, Haroldson MA (2003) Grizzly Bear. In: Feldhamer GA, Thompson BC, Chapman JA, editors. *Wild mammals of North America: biology, management and conservation*. 2nd ed. Baltimore, Maryland, U.S.A.: John Hopkins University Press. pp. 556–586.
 42. Jones ES, Heard DC, Gillingham MP, Bowman J (2006) Temporal variation in stable carbon and nitrogen isotopes of grizzly bear guardhair and underfur. *Wildl Soc Bull* 34: 1320–1325.
 43. Hellgren EC, Rogers L, Seal U (1993) Serum chemistry and hematology of black bears: physiological indices of habitat quality or seasonal patterns? *J Mammal* 304–315.
 44. Ayres KL, Booth RK, Hempelmann JA, Koski KL, Emmons CK, et al. (2012) Distinguishing the impacts of inadequate prey and vessel traffic on an endangered killer whale (*Orcinus orca*) population. *PLoS One* 7: e36842.
 45. Sapolsky RM, Romero LM, Munck AU (2000) How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21: 55–89.
 46. Wingfield JC, Hegner RE, Dufy AM, Jr., Ball GF (1990) The "Challenge Hypothesis": Theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *Am Nat* 136: 829–846.
 47. Wingfield JC, Lynn S, Soma KK (2001) Avoiding the 'costs' of testosterone: ecological bases of hormone-behavior interactions. *Brain Behav Evol* 57: 239–251.
 48. Oliveira RF (2004) Social modulation of androgens in vertebrates: mechanisms and function. *Adv Study Behav* 34: 165–239.
 49. Hirschenhauser K, Oliveira RF (2006) Social modulation of androgens in male vertebrates: meta-analyses of the challenge hypothesis. *Anim Behav* 71: 265–277.
 50. Sapolsky RM (1993) The physiology of dominance in stable versus unstable social hierarchies. In: Mason WA, Mendoza SP, editors. *Primate social conflict*. Albany, New York: State University of New York Press. pp. 171–204.
 51. Wasser SK, Papageorge S, Foley C, Brown JL (1996) Excretory fate of estradiol and progesterone in the African elephant (*Loxodonta africana*) and patterns of fecal steroid concentrations throughout the estrous cycle. *Gen Comp Endocrinol* 102: 255–262.
 52. Woods JG, Paetkau D, Lewis D, McLellan BN, Proctor M, et al. (1999) Genetic tagging of free-ranging black and brown bears. *Wildl Soc Bull*: 616–627.
 53. Kendall KC, Stetz JB, Roon DA, Waits LP, Boulanger JB, et al. (2008) Grizzly bear density in Glacier National Park, Montana. *J Wildl Manag* 72: 1693–1705.
 54. Proctor M, McLellan B, Boulanger J, Apps C, Stenhouse G, et al. (2010) Ecological investigations of grizzly bears in Canada using DNA from hair, 1995–2005: a review of methods and progress. *Ursus* 21: 169–188.
 55. Hamilton AN (2012) British Columbia Grizzly bear population estimate for 2012. Ministry of Forests, Lands and Natural Resource Operations, Victoria, BC. Available: http://www.env.gov.bc.ca/fw/wildlife/docs/Grizzly_Bear_Pop_Est_Report_Final_2012.pdf. Accessed: 30 April, 2013.
 56. Mowat G, Heard DC (2006) Major components of grizzly bear diet across North America. *Can J Zool* 84: 473–489.
 57. Meidinger D, Pojar J (1991) Ecosystems of British Columbia (Special Report Series 6, ISSN 0843-6452). Research Branch Ministry of Forests, Victoria, BC. Available: <http://www.for.gov.bc.ca/hfd/pubs/Docs/Srs/Srs06.pdf>. Accessed 30 April, 2013.
 58. Milakovic B, Parker KL, Gustine DD, Lay RJ, Walker ABD, et al. (2012) Seasonal habitat use and selection by grizzly bears in Northern British Columbia. *J Wildl Manag* 76: 170–180.
 59. Darimont CT, Price MHH, Winchester NN, Gordon-Walker J, Paquet PC (2004) Predators in natural fragments: foraging ecology of wolves in British Columbia's central and north coast archipelago. *J Biogeogr* 31: 1867–1877.
 60. Paquet PC, Alexander SM, Swan PL, Darimont CT (2006) Influence of natural landscape fragmentation and resource availability on distribution and connectivity of gray wolves (*Canis lupus*) in the archipelago of coastal British Columbia, Canada. In: Crooks KR, Sanjayan MA, editors. *Connectivity conservation*. Cambridge: Cambridge University Press. pp. 130–156.
 61. Christensen JR, MacDuffee M, Macdonald RW, Whitticar M, Ross PS (2005) Persistent organic pollutants in British Columbia grizzly bears: Consequence of divergent diets. *Environ Sci Technol* 39: 6952–6960.
 62. Paetkau D (2003) An empirical exploration of data quality in DNA-based population inventories. *Mol Ecol* 12: 1375–1387.
 63. Darimont CT, Paquet PC, Reimchen TE (2007) Stable isotopic niche predicts fitness of prey in a wolf-deer system. *Biol J Linn Soc* 90: 125–137.
 64. Darimont C, Paquet P, Reimchen T (2008) Spawning salmon disrupt trophic coupling between wolves and ungulate prey in coastal British Columbia. *BMC Ecol* 8: 14.
 65. Moore JW, Semmens BX (2008) Incorporating uncertainty and prior information into stable isotope mixing models. *Ecol Lett* 11: 470–480.
 66. Semmens BX, Moore JW (2008) MixSIR: A Bayesian stable isotope mixing model. Version 1.0. <http://www.ecologybox.org> (last accessed 11 Sept 2011). Available: <http://www.ecologybox.org>. Accessed: 11 Sept 2011.
 67. Bryan HM, Adams AG, Invik RM, Wynne-Edwards KE, Smits JE (2013) Hair is a more repeatable reflection of baseline cortisol than saliva or feces in dogs. *J Am Assoc Lab Anim Sci* 52: 189–196.
 68. BCGOV (2008) Landscape Units of British Columbia. FLNRO Resource Management Objectives Branch. Available: <https://apps.gov.bc.ca/pub/geometadata/metadataDetail.do?recordSet=ISO19115&recordUID=51078>. Accessed 30 April 2013.
 69. R Development Core Team (2011) R: A Language and Environment for Statistical Computing. Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>: R Foundation for Statistical Computing.
 70. Rohlf FJ (1975) Generalization of the gap test for the detection of multivariate outliers. *Biometrics* 31: 93–101.
 71. McCune B, Grace GB, Urban DL (2002) Analysis of ecological communities. Glendene Beach, Oregon: MjM Software Design.
 72. White DJ, Berardinelli JG, Aune KE (2005) Seasonal differences in spermatogenesis, testicular mass and serum testosterone concentrations in the grizzly bear. *Ursus* 16: 198–207.
 73. Harlow HJ, Beck TDI, Walters LM, Greenhouse SS (1990) Seasonal serum glucose, progesterone, and cortisol levels of black bears (*Ursus americanus*). *Can J Zool* 68: 183–187.
 74. Tsubota T, Nitta H, Osawa Y, Mason JI, Kita I, et al. (1994) Immunolocalization of steroidogenic enzymes P450_{sc}, 3 β HSD, P450_{c17} and P450_{arom} in the corpus luteum of the Hokkaido brown bear (*Ursus arctos yesoensis*) in relation to delayed implantation. *J Reprod Fert* 101: 557–561.
 75. Tsubota T, Takahashi Y, Kanagawa H (1987) Changes in serum progesterone levels and growth of fetuses in Hokkaido brown bears. *Bears: Their Biology and Management*: 355–358.
 76. Dye HM, Sumpter JP, Fagerlund UHM, Donaldson EM (1986) Changes in reproductive parameters during the spawning migration of pink salmon, *Oncorhynchus gorbuscha* (Walbaum). *J Fish Biol* 29: 167–176.
 77. Carruth LL, Dores RM, Maldonado TA, Norris DO, Ruth T, et al. (2000) Elevation of plasma cortisol during the spawning migration of landlocked kokanee salmon (*Oncorhynchus nerka* kennerlyi). *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* 127: 123–131.
 78. Palmer S, Nelson R, Ramsay M, Stirling I, Bahr J (1988) Annual changes in serum sex steroids in male and female black (*Ursus americanus*) and polar (*Ursus maritimus*) bears. *Biol Reprod* 38: 1044–1050.
 79. Garshelis DL, Hellgren EC (1994) Variation in reproductive biology of male black bears. *J Mammal*: 175–188.

80. Craighead J, Sumner J, Mitchell J (1995) The grizzly bears of Yellowstone. Their ecology in the Yellowstone ecosystem, 1959-92. WashingtonDC, , USA: Island Press.
81. Rode KD, Farley SD, Robbins CT (2006) Sexual dimorphism, reproductive strategy, and human activities determine resource use by brown bears. *Ecology* 87: 2636–2646.
82. Ciarniello LM, Boyce MS, Heard DC, Seip DR (2007) Components of grizzly bear habitat selection: density, habitats, roads, and mortality risk. *J Wildl Manag* 71: 1446–1457.
83. Hamilton AN (1987) Classification of coastal grizzly bear habitat for forestry interpretations and the role of food in habitat use by coastal grizzly bears. PhD Thesis. Vancouver, British Columbia, , Canada, Department of Forestry, University of British Columbia. 96p.
84. McLellan BN, Hovey FW (1995) The diet of grizzly bears in the Flathead River drainage of southeastern British Columbia. *Can J Zool* 73: 704–712.
85. Egbert AL, Stokes AW (1976) The social behaviour of brown bears on an Alaskan salmon stream. *Bears: Their Biology and Management*: 41–56.
86. Gende SM, Quinn TP (2004) The relative importance of prey density and social dominance in determining energy intake by bears feeding on Pacific salmon. *Can J Zool* 82: 75–85.
87. Albert D, Jonik R, Walsh M (1992) Hormone-dependent aggression in male and female rats: experiential, hormonal, and neural foundations. *Neurosci Biobehav Rev* 16: 177–192.
88. Ketterson ED, Nolan V, Sandell M (2005) Testosterone in females: mediator of adaptive traits, constraint on sexual dimorphism, or both? *Am Nat* 166: S85–S98.
89. Ben-David M, Titus K, Beier LVR (2004) Consumption of salmon by Alaskan brown bears: a trade-off between nutritional requirements and the risk of infanticide? *Oecologia* 138: 465–474.
90. Bellemain E, Zedrosser A, Manel S, Waits LP, Taberlet P, et al. (2006) The dilemma of female mate selection in the brown bear, a species with sexually selected infanticide. *Proc Biol Sci* 273: 283–291.
91. Donahue SW, Vaughan MR, Demers LM, Donahue HJ (2003) Serum markers of bone metabolism show bone loss in hibernating bears. *Clin Orthop Relat Res* 408: 295–301.
92. Epel E, Lapidus R, McEwen B, Brownell K (2001) Stress may add bite to appetite in women: a laboratory study of stress-induced cortisol and eating behavior. *Psychoneuroendocrinology* 26: 37–49.
93. Pravosudov VV (2003) Long-term moderate elevation of corticosterone facilitates avian food-caching behaviour and enhances spatial memory. *Proc Biol Sci* 270: 2599–2604.
94. Reneerkens J, Piersma T, Ramenofsky M (2002) An experimental test of the relationship between temporal variability of feeding opportunities and baseline levels of corticosterone in a shorebird. *J Exp Zool* 293: 81–88.
95. Pfeffer K, Fritz J, Kotrschal K (2002) Hormonal correlates of being an innovative greylag goose, *Anser anser*. *Anim Behav* 63: 687–695.
96. Herrero S (1983) Social behaviour of black bears at a garbage dump in Jasper National Park. *Bears: Their Biology and Management*: 54–70.
97. McLellan B (2011) Implications of a high-energy and low-protein diet on the body composition, fitness, and competitive abilities of black (*Ursus americanus*) and grizzly (*Ursus arctos*) bears. *Can J Zool* 89: 546–558.
98. Ebbeling CB, Swain JF, Feldman HA, Wong WW, Hachey DL, et al. (2012) Effects of dietary composition on energy expenditure during weight-loss maintenance. *JAMA* 307: 2627–2634.
99. Darimont CT, Reimchen TE (2002) Intra-hair stable isotope analysis implies seasonal shift to salmon in gray wolf diet. *Can J Zool* 80: 1638–1642.
100. Kirschbaum C, Tietze A, Skoluda N, Dettenborn L (2009) Hair as a retrospective calendar of cortisol production-Increased cortisol incorporation into hair in the third trimester of pregnancy. *Psychoneuroendocrinology* 34: 32–37.
101. Stetz J, Hunt K, Kendall KC, Wasser SK (2013) Effects of exposure, diet, and thermoregulation on fecal glucocorticoid measures in wild bears. *PLoS One* 8: e55967.
102. Koren L, Nakagawa S, Burke T, Soma KK, Wynne-Edwards KE, et al. (2011) Non-breeding feather concentrations of testosterone, corticosterone and cortisol are associated with subsequent survival in wild house sparrows. *Proc R Soc Lond B Biol Sci* 279: 1560–1566.
103. Darimont CT, Bryan HM, Carlson SM, Hocking MD, MacDuffee M, et al. (2010) Salmon for terrestrial protected areas. *Conserv Lett* 3: 379–389.
104. Levi T, Darimont CT, MacDuffee M, Mangel M, Paquet P, et al. (2012) Using grizzly bears to assess harvest-ecosystem tradeoffs in salmon fisheries. *PLoS Biol* 10: e1001303.
105. Munro RHM, Nielsen S, Price M, Stenhouse G, Boyce M (2006) Seasonal and diel patterns of grizzly bear diet and activity in west-central Alberta. *J Mammal* 87: 1112–1121.