

Agribusiness Reports

Volume 2012 *AGRIBUSINESS REPORTS* | 2012

Article 4

7-2012

The Welfare of Animals in the Veal Industry

Follow this and additional works at: <https://www.wellbeingintlstudiesrepository.org/agreports>

 Part of the [Agribusiness Commons](#), [Animal Studies Commons](#), and the [Operations and Supply Chain Management Commons](#)

Recommended Citation

(2012) "The Welfare of Animals in the Veal Industry," *Agribusiness Reports*: Vol. 2012 , Article 4.
Available at: <https://www.wellbeingintlstudiesrepository.org/agreports/vol2012/iss2012/4>

This material is brought to you for free and open access by WellBeing International. It has been accepted for inclusion by an authorized administrator of the WBI Studies Repository. For more information, please contact wbisr-info@wellbeingintl.org.





An HSUS Report: The Welfare of Animals in the Veal Industry

Abstract

Intensive confinement of calves raised for veal has long raised pointed concerns regarding the animals' welfare. Traditional production practices include individually isolating calves in narrow wooden stalls or pens, which severely restrict movement, feeding the animals an all-liquid diet deliberately low in iron, and prematurely weaning the animals. Stressful conditions lead to a high incidence of stereotypic behavior and illness. Scientific reviews of the welfare of intensively confined calves raised for veal have concluded that the young animals suffer when reared in conventional systems.

Introduction

As defined by the U.S. Department of Agriculture (USDA) Food Safety and Inspection Service (FSIS), veal “is the meat from a calf or young beef animal. A veal calf is raised until about 16-18 weeks of age, weighing up to 450 pounds [204 kg]. Male dairy calves are used in the veal industry. Dairy cows* must give birth to continue producing milk, but male dairy calves are of little or no value to the dairy farmer. A small percentage are raised to maturity and used for breeding.”¹

Two types of veal are recognized by the USDA: bob veal, produced from calves slaughtered at up to three weeks of age, and special-fed veal, from calves slaughtered around 16-18 weeks. Calves raised for both bob and special-fed veal are fed liquid milk-replacement diets.² Of the more than 450,000 calves raised for veal in the United States annually³—the vast majority of whom are male dairy calves, primarily Holsteins—approximately 15% are marketed as bob veal, with special-fed veal, also known as white (for its very pale hue), milk-fed, or formula-fed veal, comprising the bulk of the industry.⁴



Calves in veal crates.
 Photo by Farm Sanctuary.

Presently, the majority of calves raised for veal in the United States are reared indoors in individual stalls⁵ typically measuring 66-76 cm (26-30 in) wide by 168 cm (66 in) long,⁶ and approximately 30% are raised in group-housing systems.⁷ Although commonly referred to as “crates,” veal stalls are constructed with open backs and partial walls separating each individually confined calf. Calves are tied to the front of the stall with a fiber or metal 0.6-0.9 m (2-3 ft) tether,⁸ restricting virtually all movement until they reach slaughter weight.

Though not identified by the USDA in its FSIS “Veal from Farm to Table” fact sheet,⁹ a third type of veal is marketed in North America—grain-fed or non-special fed veal¹⁰—produced from calves who receive a diet that contains grain and forage (hay, silage, or pasture), as well as liquid milk replacer.^{11,12} and are typically housed

* For more information, see “An HSUS Report: The Welfare of Cows in the Dairy Industry” at www.humanesociety.org/assets/pdfs/farm/hsus-the-welfare-of-cows-in-the-dairy-industry.pdf.

in group pens or loose housing after weaning at 6-8 weeks old.¹³ The calves' flesh has a darker color¹⁴ and is often referred to as red or pink veal.

Due to strong public opinion for the improved welfare of these young animals, individual housing beyond eight weeks old has been prohibited in the United Kingdom since 1990,¹⁵ and fibrous food and bedding have been required since 2000.¹⁶ Group housing beyond eight weeks of age and daily fibrous food rations from two weeks of age have been mandatory in the rest of the European Union (EU) since January 1, 2007.^{17,18} Tethering in individual housing is not permitted and may only be used in group housing for a maximum of one hour following each milk feed.¹⁹

In the United States, a growing number of state laws prohibit or restrict the use of veal crates. Arizona voters approved a citizen initiative, Proposition 204, in November 2006, which states: "A person shall not tether or confine any pig during pregnancy or any calf raised for veal, on a farm, for all or the majority of any day, in a manner that prevents such animal from lying down and fully extending his or her limbs, or turning around freely."²⁰ The law went into effect December 31, 2012, and prohibits the housing of calves in individual stalls. Colorado followed suit in 2008, banning crates for veal calves effective January 1 2012 and gestation crates for pregnant pigs by 2018.^{21,22} Veal crates were also banned by the landslide passage of Proposition 2 on the November 2008 California ballot. The measure passed with 63.5% of the vote, when more than 8 million Californians elected to disallow the intensive confinement systems of veal crates, gestation crates for pregnant sows, and battery cages for egg-laying hens, effective January 1, 2015.^{23,24,25} In May 2009, Maine passed a law banning veal crates for calves and gestation crates for sows, effective January 1, 2011.²⁶ Michigan followed in October 2009, with passage of state legislation to phase out veal crates by 2012 and gestation crates within ten years.²⁷ In 2010, the Ohio Livestock Care Standards Board was created,²⁸ and the Board issued a rule effective January 1, 2018 that requires calves to be group housed by ten weeks of age.²⁹ In June of 2012, Rhode Island passed legislation banning both veal and gestation crates.³⁰

In 2007, the American Veal Association's board of directors unanimously approved a policy that the veal industry fully transition to group housing production by the end of 2017. As reported by the industry journal *Feedstuffs*, this policy resolution was due in part to the fact that "[v]eal customers and consumers are concerned with current individual stall systems, and how animals are raised is increasingly part of customer and consumer purchasing decisions."³¹ Progress toward this goal has been substantial. In 2012, an American Veal Association (AVA) survey found that 70 percent of veal calves raised by AVA members would be group housed by the end of the year.³²

In January 2007, two of the nation's largest veal producers, Strauss Veal³³ and Marcho Farms,³⁴ both announced corporate policies to convert their operations to crate-free group housing within 2-3 years due to animal welfare concerns. As reported by *Meat Processing*, Strauss Veal & Lamb International "is committed to raising veal calves in a more humane manner. The company's goal is to be 100-percent converted to raising calves by the European-style, group-raised method within the next two to three years." Randy Strauss, co-president and CEO, stated to the industry journal that "this is the right thing to do...The traditional way of raising veal calves involves putting each calf in an individual stall. This practice is increasingly being frowned upon by a growing number of customers and consumers alike throughout the world."³⁵ Strauss Veal & Lamb is now free of crates and tethers.^{36,37} By 2010, 50% of calves at Marcho Farms were kept in some sort of loose housing arrangement.³⁸

After a comprehensive two-year study, the independent Pew Commission on Industrial Farm Animal Production, a project of The Pew Charitable Trusts and the Johns Hopkins Bloomberg School of Public Health chaired by former Kansas Governor John Carlin and including former U.S. Agriculture Secretary Dan Glickman, concluded that veal crates should be phased out:

After reviewing the literature, visiting production facilities, and listening to producers themselves, the Commission believes that the most intensive confinement systems, such as restrictive veal crates, hog

gestation pens, restrictive farrowing crates, and battery cages for poultry, all prevent the animal from a normal range of movement and constitute inhumane treatment.³⁹

Despite important movement toward the group-housing of calves raised for veal by U.S. and European industries, legislative bodies, and consumers, individual housing and restrictive tethering of calves are not the only aspects of veal production that degrade calf welfare. Many customary veal industry practices fail serious scientific examination of their impacts on calf health and well-being.

Indeed, research has shown that many of the welfare problems associated with customary U.S. veal industry practices could be substantially reduced or eliminated by group housing on straw, teat-feeding, and the provision of solid feed. In a study on the effects of different types of husbandry on calf behavior, Webster *et al.* concluded that “the practice of rearing calves for veal on teats and in groups appears to be ‘natural’ in the sense that their behavior is, in many respects, similar to that of calves at pasture with their mothers.”⁴⁰ Andrighetto *et al.* conducted a similar study and determined: “Group pen calves had the opportunity for locomotion and social behaviour and were allowed to adopt more comfortable resting postures. The improved welfare of calves kept in group pens was confirmed by the higher haemoglobin levels at the end of the growing cycle.”⁴¹ Sabbioni *et al.* found that group-housed calves had higher growth performance and better food conversion ratios than individually housed calves. They concluded that the lower incidence of stress in the group housed calves, as confirmed by lower cortisol levels, may be the reason for their lower energy consumption.⁴²

As well, in its *Report on the Welfare of Calves*, the European Commission’s Scientific Veterinary Committee (SVC) wrote that “general comparisons indicate that the housing of calves in individual pens, and the tethering of calves, result in problems for their welfare which are significantly reduced when the calves are group-housed on straw.”⁴³ In a study on the welfare impacts of solid feed supplements for veal calves, Morisse *et al.* concluded that supplying calves with straw-cereal pellets helped reduce non-nutritive chewing and was “positive for the physiological aspects of welfare.”⁴⁴ Cozzi *et al.* found that providing solid feed to veal calves promoted forestomach development and improved calves’ health status, with fewer iron treatments for anemia and medical treatments for respiratory and gastrointestinal diseases necessary.⁴⁵

In sum, a review of the welfare of calves in veal units reported: “Although calves are sucklings of social-living ruminants, veal calves are not allowed to suck, to have a social life or to ruminate. This, added to anaemia in order to obtain white meat, and the high rate of morbidity caused by high density of the animals, sums up the welfare problems which arise.”⁴⁶

Impacts of Unnaturally Early Weaning

Cows, like all mammals, produce milk to feed their young and, as such, must give birth to a calf for lactation to begin. On typical commercial dairy farms, cows are impregnated and calve on a yearly cycle for milk production, and calves are usually separated from their dams within a few hours of birth.⁴⁷ Female calves to be used as replacement dairy cows are raised in individual pens and fed milk replacer until 4-12 weeks old when they are moved to group pens and weaned onto solid food.⁴⁸ As male calves are often not desired breeds for beef production, they are commonly sold to the veal industry, primarily through livestock auctions.⁴⁹ According to the American Veal Association, “The special fed/milk fed veal industry was born in the United States in the mid 1960’s [*sic*] in part, as a result of a need to relieve the US dairy industry of its unneeded byproducts.”⁵⁰

In natural cattle herds, dam and calf forge very strong and long-lasting bonds. Reinhardt and Reinhardt studied cow-calf relationships in a herd of semi-wild Zebu cattle (*Bos indicus*) over five years and found that “all cows preferred their daughters and sons over non-related calves as licking and as grazing partners for several years.” The birth of a second, third, or even fourth calf failed to disrupt the close association between the cow and her older offspring. Female calves remained attached to their dams even after having themselves calved once or twice, and relationships between cows and their male offspring were evident after four and one-half years, past the son’s sexual maturity.⁵¹ Vessier *et al.* observed similar behavior in a herd of domestic beef cattle (*Bos taurus*). The cow and her yearling calf had more contacts and stayed closer to each other than did unrelated

cows and yearlings, and that close bond remained unbroken even after she gave birth to a new calf. The presence of the yearling did not alter the establishment of the maternal bond with the newborn calf.⁵²

Maternal Deprivation

The routine early separation of cows and their calves in the dairy and veal industries is distressing for both.⁵³ Hudson and Mullord found “that 5-min contact with a calf immediately post partum is sufficient for the formation of a strong, specific maternal bond with that calf.”⁵⁴ Calves separated from their dams at birth, as observed by Lidfors,⁵⁵ were less active and vocalized and licked themselves more than calves remaining with their mothers. Marchant-Forde *et al.* reported that calves separated from their dams 24 hours after birth recognized and responded to recordings of their dams’ calls 24 hours after separation, with the cows’ vocalizations eliciting cardiac and behavioral responses in their calves.⁵⁶

In its 1995 *Report on the Welfare of Calves*, the SVC concluded: “The best conditions for rearing young calves involve leaving the calf with the mother in a circumstance where the calf can suckle and can subsequently graze and interact with other calves.”⁵⁷

Colostrum Deficiency and Impaired Immune Function

Newborn calves have no antibodies against infections and are entirely dependent on immunoglobulin in mother’s milk for immunological protection.⁵⁸ Colostrum, the milk dams produce during the first few days after calving, is especially high in immunoglobulin. Adequate intake of colostrum is critical for the future health of the calf, as those with low concentrations of absorbed immunoglobulin are more susceptible to diarrhea.⁵⁹ The routine practice of removing newborn calves from their dams within a few hours of birth may jeopardize this important transfer of immunoglobulin.

Although colostrum is collected from recently calved cows, in top dairy-producing states such as California, it is often sold to facilities specializing in raising female calves for the dairy industry,⁶⁰ as their long-term health is considered to be of greater value than that of male calves reared for veal. Surveys of U.S. veal farms confirm that many calves do not receive adequate colostrum. Stull and McMartin, in a study of western U.S. veal facilities, reportedly found that only one in five (20%) of calves entering veal units had received adequate colostrum.⁶¹ Stull and McDonough reported similar results after their evaluation of the welfare of calves at ten commercial U.S. veal farms, determining that only 22% had received adequate transfer of colostral immunoglobulins, leading them to conclude that “[t]he major factor likely to adversely affect the welfare of the veal calf was an inadequate immune system on its arrival at the veal facility.”⁶²

The colostrum delivery method is important as well. Lidfors found that calves who receive colostrum by suckling their dams have lower mortality rates and higher serum immunoglobulin concentrations than those given colostrum from an open bucket.⁶³ Weary and Chua found that calves kept with their dams for four days after birth had fewer bouts of diarrhea during the first three weeks of life than calves separated at six hours or one day, despite the fact that all the calves were bottle-fed colostrum within 24 hours of birth.⁶⁴ Metz and Metz showed that dam-reared calves defecate and urinate earlier after birth than artificially raised calves, likely as a result of frequent licking of the calf by the dam, and concluded that the early removal of the meconium (first excretion of feces) by the dam promoted colostrum intake and digestive functions in the calf.⁶⁵ However, calves of highly productive dairy cows are not always able to ingest adequate amounts of colostrum to keep them healthy through natural nursing.^{66,67,68,69} The reasons for this include ineffective or delayed nursing caused by mastitis (a common bacterial infection of the udder) or other illness of the dairy cow, and her prevailing physical characteristics such as low pendulous udders and large teats.⁷⁰

Denial of Natural Sucking Behavior

On most U.S. commercial veal farms, calves are bucket-fed liquid milk replacer twice per day, in contrast to the four to ten daily suckling sessions they would naturally have with their dams.⁷¹ According to Donald Broom,

Colleen Macleod Professor of Animal Welfare at the University of Cambridge Department of Clinical Veterinary Medicine, the biological mechanisms that have evolved in order for calves to receive nutrients are searching behaviors, which should result in finding a teat, followed by licking and sucking to ingest milk. These behaviors are not eliminated if the gut is filled with milk—that is, the licking and sucking behavior itself is greatly important to the calf.⁷² Hammell *et al.* found that calves who were bucket-fed milk replacer *ad libitum* sucked an artificial, dummy teat that did not supply milk for 13 minutes a day and that the sucking of the dummy teat often interrupted milk-drinking. The researchers concluded that “[t]his study underlines the strong need for sucking in the young calf. Sucking is maintained even while not rewarded by milk, and it can even interrupt milk drinking. The response is not even cancelled by satiation with milk.”⁷³ Sucking is also important for the release of metabolic, gastrointestinal hormones that aid the digestive process and are thought important for satiety.⁷⁴

In young mammals, the motivation to suck is assumed to be strong, as under natural conditions their survival would depend on it. Denial of this important, instinctive behavior by bucket feeding, for example, often results in non-nutritive sucking of such objects as the enclosure or tether, or of other calves,⁷⁵ known as cross-sucking. Cross-sucking is typically directed at the ears, mouth, navel, scrotum, and prepuce, and occurs primarily within 10-15 minutes after milk feeding.⁷⁶ Cross-sucking, which is believed to increase disease transmission,⁷⁷ has never been reported in calves raised by their dams or nurse cows.⁷⁸ Tethering is a common intervention used to prevent cross sucking. Numerous studies, however, have shown that providing calves with an artificial teat to satisfy their need to suck, as well as fibrous foods such as hay, largely eliminates cross-sucking in group-housed calves.

In a 2002 study published in the *Journal of Animal Science*, the behavior of bucket-fed calves was compared with those fed from an artificial teat.⁷⁹ During the feeding, bucket-fed calves had higher heart rates than teat-fed calves and, after the meal, engaged in non-nutritive sucking of the bars, themselves, and their conspecifics. Teat-fed calves took longer to consume the milk replacer and laid down more quickly after the meal. The researchers concluded that “teat-feeding reduces non-nutritive oral activities after the meal and induces a calmer state [in the calves] than bucket-feeding.”⁸⁰ Another study found that feeding calves through an artificial teat with a small orifice (to slow flow rate and increase sucking time), combined with the provision of hay at the end of a milk meal, greatly reduced cross-sucking in group-housed calves.⁸¹ These results echo previous findings that providing milk through an artificial teat as well as supplying hay significantly reduces non-nutritive sucking after a milk meal.⁸² Free access to water around meals can also help reduce the occurrence of nonnutritive oral behavior.⁸³

In the EU, where group housing of calves raised for veal is required after eight weeks of age,⁸⁴ tethering of group-housed calves for one hour following each milk meal is permitted to help reduce cross-sucking.⁸⁵ However, a 2003 review of the effects of feeding method on cross-sucking in group-housed calves concluded: “[T]he best way of preventing cross-sucking is to provide an outlet for the natural motivation to suck in connection with ingestion of milk...From an animal welfare point of view, the use of milk feeding methods that provides [*sic*] an outlet for the motivation to suck is preferred to tethering after the milk meal. Firstly, because a tethered animal may feel frustrated if they are highly motivated, but not able, to suck....Moreover, sucking in itself has beneficial physiological consequences for the calf in relation to digestion.”⁸⁶

Impacts of Diet

Cattle are ruminants, animals who digest their food in two stages. They have evolved to efficiently digest grass and other fibrous plants by chewing and swallowing once, regurgitating later, and chewing again at their leisure,⁸⁷ a practice commonly known as “chewing the cud.” Calves raised outdoors begin to eat grass and ruminate after a few weeks of age; by four months, the young animals may spend six hours a day grazing and several hours ruminating.⁸⁸

In contrast, the majority of calves in the U.S. veal industry are fed only liquid milk replacer.⁸⁹ This unnatural diet, low in fiber, is also formulated to be low in iron to pale the calves’ flesh.⁹⁰ Consumers are generally

believed by the veal industry and the retail sector to assess veal quality on color, with whiter meat commanding higher prices.⁹¹ As the color of the calves' flesh is greatly influenced by the amount of iron in the diet, feeding animals exclusively milk replacer achieves the desired minimal iron intake.⁹²

Feeding calves all-liquid diets limits the physiological development of the forestomach and the normal process of rumination. Cozzi *et al.* compared the growth performance and forestomach development of calves fed only milk replacer with calves fed the same liquid diet plus 250 g (8.82 oz) of dried beet pulp or wheat straw. In calves provided solid feed in addition to milk replacer, a clear progression in forestomach development was observed, as well as a marked reduction in the number of hairballs.⁹³ The finding is particularly noteworthy as large hairballs can clog the rumen, resulting in digestive problems and even death.⁹⁴ Morrissette *et al.* studied changes in the rumen of calves fed either an all-liquid diet or ones supplemented with 10-25 kg (22-55 lb) of pelleted straw and cereals. Compared with calves on the all-liquid diet, both groups of calves on pellet-supplemented diets showed increased reticulo-rumen weight, the presence of small papillae that help nutrient absorption from food, and significantly fewer hairballs.⁹⁵

The SVC recommends: "Every calf should be fed a daily source of long fibre to stimulate the development of villi in the rumen, and the long fibre should be supplemented with a fermentable material such as starch to maintain the microbial flora of the gut. They should receive a minimum of 100 g [3.53 oz] of roughage per day from 2 to 15 weeks of age, increasing to 250 g [8.82 oz] per day from 15 to 26 weeks of age but it would be better if these amounts were doubled."⁹⁶ This recommendation forms the legal minimum requirement for fibrous food for calves raised in the United Kingdom,⁹⁷ while for the remaining European Union member countries, the legal minimum requirement at 2 weeks of age is slightly lower, at 50 g (1.76 oz) per day.⁹⁸

Iron Deficiency

The already low iron content of all-liquid milk-replacer diets commonly fed to calves raised for veal is reduced at the beginning of the fattening period.⁹⁹ A deficiency in iron can inhibit the body's ability to produce red blood cells and result in reduced hemoglobin concentrations and eventually iron-deficiency anemia. Stull and McDonough's evaluation of calf welfare at ten commercial U.S. veal units found that one in four animals was marginally anemic (blood hemoglobin concentration under 4.9 mmol/l) and one in ten was clinically anemic (blood hemoglobin concentration under 4.34 mmol/l) when sent for slaughter.¹⁰⁰

According to the SVC's *Report on the Welfare of Calves*, many studies "indicate impaired performance and an increased disease susceptibility in calves whose blood haemoglobin concentration is below 4.5 mmol/l....It is therefore recommended that calves should be fed sufficient dietary iron to maintain the haemoglobin concentration at a minimum of 4.5 mmol/l until slaughter."¹⁰¹ In 2006, the European Food Safety Authority (EFSA) recommended blood hemoglobin concentrations to be maintained at a minimum of 6.0 mmol/l throughout the life of the calf,¹⁰² although the European Union legal minimum remains at 4.5 mmol/l.¹⁰³

Numerous scientific studies have linked milk-replacer diets with insufficient iron levels. For example, Welchman *et al.* compared blood hemoglobin concentrations in calves fed milk replacer either alone or with solid feed of varying iron content and found that calves raised exclusively on milk replacer had levels indicating iron-deficiency anemia at 16-20 weeks; only calves who received solid feed with substantial iron content showed hemoglobin concentrations similar to calves raised for beef production.¹⁰⁴ Reece and Hotchkiss also reported the development of iron-deficiency anemia in calves fed milk replacer only, both in individual housing and group housing, compared to an increase in hemoglobin concentration over 15 weeks in calves fed hay and grain.¹⁰⁵ Similarly, the study by Morrissette *et al.* found that calves on milk-replacer diets supplemented with either 10-25 kg (22-55 lb) of solid food had higher hemoglobin concentrations than those fed exclusively milk replacer; however, only those calves receiving the supplemental 25 kg (55 lb) of solid food were in conformity with the EU minimum requirement.¹⁰⁶ In Cozzi *et al.*'s investigation, calves receiving milk replacer plus 250 g (8.82 oz) of dried beet pulp or wheat straw per day required fewer iron treatments for anemia than those fed milk replacer only and needed fewer medical treatments for respiratory or gastrointestinal diseases, and only those calves fed beet pulp had a darker carcass at slaughter, not those given wheat straw.¹⁰⁷

Inactivity may also contribute to iron deficiency, which is of particular concern for individually housed calves who are typically tethered and severely restricted of movement. Although both group- and individually housed calves on milk replacer-only diets developed anemia in Reece and Hotchkiss' study, those animals in group housing had higher hemoglobin concentrations than calves in stalls.¹⁰⁸ Xiccato *et al.* also found that group-housed calves had higher hemoglobin concentrations than individually housed calves even though both groups of calves received the same diet.¹⁰⁹ The authors of both studies concluded that the higher physical activity of group-housed calves led to a muscle oxygen deficit that stimulated red blood cell production, whereas the persistent inactivity of individually housed calves did not stimulate red blood cell production as the muscles were not oxygen-deprived.¹¹⁰

Stereotypic Behavior

On all-liquid diets, calves cannot perform the normal behaviors of rumination and chewing. The natural instinct of calves to ruminate and manipulate material with their mouths is so strong that milk-fed calves will perform sham activities, attempting to ruminate despite not having ingested roughage and engaging in vacuous oral motions.¹¹¹ As a result, calves develop oral activities such as tongue-rolling and sucking, licking, and biting inanimate objects.¹¹² Social deprivation,¹¹³ the inability to groom the hindquarters,¹¹⁴ and the inability to explore¹¹⁵ can exacerbate purposeless oral actions in individually housed calves. Purposeless oral behavior is identified by animal welfare expert Donald Broom as “abnormal behavior that is not shown by calves that can cope well with their environment.” “[J]ust as in man, zoo animals or sows,” Broom has written, “stereotypies are indicators of poor welfare.”¹¹⁶

The provision of solid feed, particularly hay, reduces the frequency and incidence of some non-nutritive oral activities such as chewing.¹¹⁷ Mattiello *et al.* found purposeless oral behaviors were more common in calves fed only milk replacer and lowest in calves also fed straw. By the end of the fattening period, however, the addition of 250 g (8.82 oz) of straw per day, the EU legal minimum, no longer prevented abnormal oral behavior as this amount was presumably unable to satisfy the needs of growing calves. The researchers concluded that 250 g (8.82 oz) per day of fibrous food for calves aged 15-20 weeks is inadequate to reduce abnormal oral behavior at the end of the fattening period.¹¹⁸

Vessier *et al.* also found that provision of solid feed largely reduced the amount of non-nutritive nibbling performed by calves. Tongue-rolling, however, seemed to come from a cumulative effect of lack of roughage and social deprivation, since it occurred mainly in calves both housed in individual stalls and fed milk replacer.¹¹⁹ Webster *et al.* identified a similar cumulative effect of social isolation and lack of roughage when studying different rearing systems on calf behavior. Calves in individual stalls without solid food spent 16% of their time in purposeless oral activity, while the mean for all other groups, including those group-housed without solid food, was 2%.¹²⁰

Abomasal Damage

Stress in mammals is often reflected in stomach-wall damage, and calves raised for veal frequently show damage of the abomasal wall (the lining of the fourth compartment of the stomach).¹²¹ Differences in occurrence and degree of ulceration may be related to coping capabilities among calves.¹²² Stereotypic behavior such as tongue-rolling may help individually housed calves cope in an environment that lacks social and environmental stimulation by increasing sensory stimulation and muscular and skeletal activity, and giving them a sense of control over their relationship to the environment.¹²³

Wiepkema *et al.* studied the relationship between behavior and abomasal damage in calves. In their study, all of the calves were housed in individual crates and bucket-fed milk replacer twice daily with no solid food supplements. At slaughter, 67% showed abomasal damage, either ulcers, scars, or both. However, those calves who had developed tongue-rolling (33% of the animals in the study) had no ulcers or scars, while all of the calves who had not developed tongue-rolling (67% of the animals) did have ulcers or scars. The occurrence of

biting or licking had no relationship to abomasal damage, suggesting that these abnormal oral behaviors are more related to other factors, such as lack of solid feed, than to stress.¹²⁴

While coarse straw as a feed supplement has been shown to enhance abomasal damage, calves fed milk replacer only also develop this damage.¹²⁵ According to the SVC, when calves raised for veal receive a solid feed supplement that is balanced for starch, fiber, and degradable protein, the amount of abomasal damage is less than in conventional all-liquid diets.¹²⁶

Impacts of Intensive Confinement

Presently, the most common U.S. veal production housing system is the open-backed stall in which calves are separated by wooden, slatted partitions and individually tethered around the neck by a 0.6-0.9 m (2-3 ft) chain or rope affixed to the front of the enclosure.¹²⁷ In its 1995 report, the SVC concluded:

The welfare of calves is very poor when they are kept in small individual pens with insufficient room for comfortable lying, no direct social contact and no bedding or other material to manipulate....Every calf should be able to groom itself properly, turn around, stand up and lie down normally and lie with its legs stretched out if it wishes to do so.¹²⁸

Neither the open-backed stall nor the individual pen permit all of these behavior patterns.

Restriction of Movement

Among the greatest deprivations individually housed calves suffer are their ability to adopt their preferred lying posture and to stand and lie down naturally.¹²⁹ As a primary purpose of lying down is to relax certain muscles, the restrictions that stalls and tethers place on most normal lying postures of calves may impede full relaxation of the body and prevent the animals from lying comfortably.¹³⁰ For all young mammals, rest is critical, and sleep disruption may occur if certain lying positions cannot be adopted.¹³¹ Lying posture is also very important for thermoregulation. Stretching out the legs laterally can help maximize surface area to prevent overheating.¹³²

Several studies have shown that calves tethered in stalls or confined in narrow pens cannot adopt the postures they would choose for most effective body temperature regulation or comfort. In Stull and McDonough's evaluation of the welfare of calves in ten commercial U.S. veal units, nine facilities used stall and tether systems and one used group housing. The researchers found that group-housed calves extended one or more legs when lying down 13% of the total lying time, whereas calves in stalls extended their legs only 2% of the lying time.¹³³ These results were echoed by Andrighetto *et al.* who found that individually housed calves spent more time lying with all four legs bent, while group-housed calves more frequently adopted postures with one or more legs outstretched.¹³⁴

Smits and de Wilt found that the incidence of lying with the head turned back on the shoulder or belly decreased more rapidly over time for calves in individual pens than for those group-housed, and the occurrence of lying with the head reclined forward on the floor increased over time in the individual, but not group, pens. The researchers concluded that lying with the head turned backwards is thwarted by the sides of the pen.¹³⁵ Similarly, Webster *et al.* determined that compared with those in group pens, calves in individual pens spent less time sleeping at two-weeks-old and concluded that their inability to adopt the normal sleeping posture with the head tucked into the flank may be a factor.¹³⁶

Both individual pens and open-backed stalls used with tethers restrict movement. Wilson *et al.* compared the stress and behavioral effects of enclosed pens without tethers and individual partial stalls with tethers. Both pens and stalls measured 179 cm (70.5 in) in length and came in three different widths: 56, 66, and 76 cm (22, 26, and 30 in). They found no consistent differences in postures or behavior among calves in the two individual housing systems, however calves housed in the narrowest pens 56 cm (22 in) were unable to extend their legs while recumbent, and had difficulty changing from a lying to a standing position. Tongue-rolling and

purposeless chewing were exhibited in both stall and pen treatments, as well as increased cortisol levels over time, which suggests an increased degree of stress in individual housing designs regardless of the use of tethers. The researchers concluded: “Few consistent differences were observed in stress indicators...in special-fed veal calves as a result of tethering in partially divided stalls vs nontethering in enclosed individual pens.”¹³⁷

Tether and stall systems are no more welfare-friendly than individual pens, and several studies have shown them to be similarly poor for calf welfare. Van Putten compared the behavior of 150 calves housed in the same building but kept in three different housing systems: individual crates, individually tethered between partitions, or in groups of three. No significant differences in measured behavioral patterns were found between the crated and tethered calves. A tendency towards greater grooming frequency noted for the tethered calves was explained by the fact that the observed grooming behaviors were often only failed attempts at grooming aborted due to the shortness of the chain around their necks. The authors concluded: “Tethering is not a real solution [from a welfare perspective] because of the restriction of grooming and lying down.”¹³⁸

Le Neindre *et al.* also compared the behavior of individually tethered versus group-housed calves and found that those who were tethered had difficulty stretching their legs and interacted less with other calves. The researchers concluded: “Tethering for the entire resting period is not a good system for maintaining a minimum acceptable standard of welfare for veal calves because it limits their activities.”¹³⁹

The 2006 *Scientific Report on the Risks of Poor Welfare in Intensive Calf Farming Systems* by the EFSA (an update to the SVC’s 1995 *Report on the Welfare of Calves*) echoed this conclusion, stating: “Tethering always causes problems for calves. Calves housed in groups should not be tethered except for periods of not more than one hour at the time of the feeding of milk or milk substitute. Individually housed calves should not be tethered.”¹⁴⁰ This opinion was enshrined in EU law and went into effect on January 1, 2007.¹⁴¹

Unsuitable Flooring in Open-Backed Stalls and Individual Pens

Flooring type has influence over calf movements, particularly getting up and down, lying, and resting. Most calves raised for veal in the United States are kept on bare wooden slats or a plastic-coated metal grating,¹⁴² while under U.K. law, calves must be provided “appropriate bedding” from birth until slaughter.¹⁴³ When cattle are allowed to choose between different floor types, they reportedly prefer deep litter to slatted floors, especially for resting,¹⁴⁴ indicating welfare would likely be improved with the provision of bedding over barren flooring.

Wooden slatted floors absorb liquid from manure and become slippery,¹⁴⁵ making it difficult for calves to get up and down,¹⁴⁶ which can lead to leg disorders. According to the SVC’s report, studies have found that calves raised for veal who were housed on wooden slatted floors showed damage of the carpal joints at slaughter, whereas those kept on rubber flooring slipped less and had less damage to their carpal joint.¹⁴⁷

Given the importance of adequate rest for growing animals, a suitable floor is particularly important for calves, particularly those raised in barren enclosures without appropriate sleeping and resting quarters. Webster *et al.* compared the behavior of calves in several husbandry systems, including at pasture with their dams, group-housed in straw yards, in individual pens on concrete floors, and in individual crates with slatted wooden floors. The researchers found that, compared to calves in any other housing regimen, those in wooden crates on slatted floors spent less time lying down and almost twice as long standing idle, often appearing to be standing insecurely, suggesting they were reluctant to change positions possibly because they found the surfaces slippery.¹⁴⁸ Indeed, reported the SVC, slippery slatted floors may result in decreased frequency of lying bouts.¹⁴⁹

Lack of Exercise

Calves, like all young mammals, have evolved with a need for regular exercise, which helps reduce problems associated with inactivity,¹⁵⁰ such as abnormal bone and muscle development and joint disorders. Intensive confinement systems prohibit exercise and normal muscle growth in order to produce tender veal.¹⁵¹

If healthy calves are raised in an environment with ample space to play vigorously, they will gallop, buck, and kick.¹⁵² If housed with other calves, they will also engage in play fighting.¹⁵³ In contrast, when closely confined for prolonged periods, these normal behaviors are frustrated, resulting in an intensification of the drive to perform these activities.¹⁵⁴

Dantzer *et al.* compared the behaviors of calves tethered in stalls with group-housed calves when released into an open field. Tethered calves exhibited higher activity scores and spent less time immobile, suggesting a compensatory reaction to deprivation of these activities when restrained.¹⁵⁵ Similarly, Dellmeier *et al.* found that calves housed in individual stalls or pens for six weeks exhibited more locomotive and social behavior when released in an open field than calves housed in groups. Only stall- and pen-confined calves stumbled and fell during the field test, suggesting that prolonged inactivity had inhibited muscle development and coordination. The researchers concluded that “[t]he greater incidence of highly active behaviors observed for the more confined calves during open field testing, despite their tendency to stumble and fall, attests to the strength of motivation induced by ethostasis of locomotion.”¹⁵⁶

Social Deprivation

Cattle are social animals who obtain physical, physiological, and psychological comfort from each other.¹⁵⁷ Under natural conditions, calves from two weeks of age start to associate in groups during the day while their mothers forage and begin to form relationships with their peers.¹⁵⁸ In Reinhardt and Reinhardt’s study of semi-wild Zebu cattle, the majority of calves formed close relationships with one or two other non-related calves of a similar age and maintained those bonds over three or more years. The scientists concluded: “In natural cattle herds the social structure is based on matriarchal families which in their turn are interconnected by means of friendship relationships between non-kin partners.”¹⁵⁹ For calves raised without their mothers, social contact with other calves is particularly important.¹⁶⁰

Tethering calves in open-backed stalls prevents the animals from adequate social contact. Holm *et al.* investigated calves’ motivation for access to full social contact as compared with head-only contact. The researchers found that while calves were willing to work to gain access to both types of contact, they were more motivated to gain access to full social contact than head-only contact and concluded that “calves’ welfare may be threatened if they are not allowed to perform social behaviours, and since motivation is apparently higher for full social contact than for head contact it is likely that their welfare will be better if housed in groups....”¹⁶¹

When calves are confined, their motivation for social contact intensifies. Dellmeier *et al.* studied the effect of four methods of confinement on calf behavior: tethered in individual stalls, individually penned, in outdoor hutches open at one end and restrained on a 2.5 m (2.7 yd) chain, and group housed in an outdoor yard. After six weeks, the calves were individually tested in an open field in the presence of unfamiliar calves. According to the authors: “The number of social encounters in which calves engaged increased with increasing degree of confinement....Social encounters were typically initiated by licking the ears, face, or neck of another calf, rubbing foreheads together, or butting heads.”¹⁶²

Given the natural behaviors of calves, the EFSA concluded: “Since calves are social animals they should be kept in social groups wherever possible.”¹⁶³

Inability to Explore

All animals explore their surroundings in part to discover sources of danger as well as areas of retreat or escape.¹⁶⁴ Calves reared on pasture spend considerable time investigating their environment through visual, auditory, tactile, olfactory, and gustatory cues.¹⁶⁵ When tethered in stalls, however, their ability to explore is severely restricted, limited primarily to sniffing and licking the front part of the enclosure.¹⁶⁶ The negative impacts of denial of exploratory behavior on calf welfare were outlined in the SVC’s 1995 report:

Calves given little space, low environmental complexity and no variety in their environment have little possibility for exploration and this may result in poor welfare as indicated especially by high levels of abnormal behaviour....It is likely that the inability to explore, and to escape from perceived danger contributes to the high level of oral stereotypies, self licking and hair ingestion which occurs in calves confined in individual pens.¹⁶⁷

In addition, the inability to investigate surroundings can lead to increased fearfulness in individually housed calves. de Wilt reportedly observed the behavior of calves raised for veal in various housing systems and concluded that those animals tethered in individual stalls are more easily alarmed because their ability to explore their environment is significantly restricted.¹⁶⁸ Similarly, Jensen *et al.* found that calves reared in individual stalls for prolonged periods are more fearful than group-housed calves. They housed calves in four different types of housing (small single pen, large single pen, small group pen, and large group pen) for three months, twice as long as Dellemeier *et al.*, and then introduced them to a novel environment. In open-field tests with an unfamiliar calf present, individually reared calves from both small and large pens were reluctant to sniff the calf and had high heart rates, and a greater percentage of individually reared calves were afraid to enter the arena than group-reared calves.¹⁶⁹ These findings echo the EFSA's 2006 report that stated: "Calves need to explore and it may be that higher levels of stereotypes and fearfulness in poorly lit buildings or otherwise inadequate conditions are a consequence of their inability to explore."¹⁷⁰

Inability to Groom

In order to maintain hygiene and help prevent disease, animals groom themselves when necessary, principally by licking themselves. Stimuli to groom may emanate from anywhere on the body, and once grooming is initiated, there is an urge to groom the whole body.¹⁷¹ Cattle naturally lick all the accessible parts of their bodies, including their hind limbs and tails, and will often rub against objects such as branches and fence posts to reach inaccessible parts.¹⁷² Calves tethered to the front of stalls, however, are unable to rub the hind parts of their body as the design of the open-backed stall lacks objects or structures behind the animals. Licking of the hindquarters is also greatly restricted by the tether and sides of the stall. Excessive licking of the forelegs, a re-directed behavior, is common in stall and tether systems.¹⁷³

Stress

Individual pens and stall and tether systems do not permit calves to respond to many types of stimuli, both internal, such as the need to scratch an itch, and external, such as the need to avoid a perceived threat, and, as discussed above, also frustrate many of the calves' drive-motivated behaviors, including grooming, exploring, and interacting socially. The chronic deprivation of these needs and behaviors can lead to stress.¹⁷⁴

Dantzer *et al.* showed that calves tethered in stalls had higher cortisol responses to adrenocorticotrophic hormone (ACTH) than group-housed calves, demonstrating that the adrenal glands of tethered calves were more active than those of loose calves, a physiological indicator of stress.¹⁷⁵ Similarly, Friend *et al.* found that calves tethered in stalls had higher adrenal responses to ACTH than group-housed calves, as well as increased levels of thyroid hormones and a higher neutrophil to lymphocyte ratio, another physiological indicator of chronic stress.¹⁷⁶ Raussi *et al.* found that the presence of another calf greatly reduced the stress of confinement for calves in the veal industry. In their study, individually housed calves who could see, sniff, touch, and lick calves in adjacent pens through open wooden partitions were compared to calves housed in pairs in larger pens. The individually housed calves had higher cortisol responses to ACTH than pair-housed calves. The scientists concluded that calves feel a need for social contact and that pair-housing can reduce the stress due to separation from conspecifics.¹⁷⁷

Disease

Young calves are susceptible to pathogens and individual housing is used, in part, because it may help to reduce the transmission of pathogenic organisms by minimizing animal-to-animal contact.¹⁷⁸ Yet, according to Friend and Dellmeier, who examined the problems relating to artificially rearing veal calves:

[V]irtually all crates are wooden and can readily harbor microorganisms, and the vast majority of them have slatted partitions separating calves, which would not inhibit transmission of airborne organisms. Typically calves can also touch each other's head and frequently do make oral contact and nasal contact...possibly because this is one of the few sources of stimulation available to them... Thus, there remain many avenues for transmission of disease in most crate-housing systems.¹⁷⁹

Diarrhea and pneumonia are common diseases in commercial veal production facilities.¹⁸⁰ The stresses of transport to the livestock auction and then the veal unit, the mixing of calves from different sources, the possibility of immune compromise as a result of inadequate immunoglobulin transfer, nutritional inadequacies of an all-liquid diet, and over-intensive stocking can predispose calves to pathogens.¹⁸¹ In a study of eight commercial veal units in the United States, McDonough *et al.* found that 92% of ill calves had not received sufficient transfer of passive immunity from their dams. Of those calves who died from diarrhea, all had had complete failure of immunoglobulin transfer.¹⁸²

McFarlane *et al.* studied the effect of iron intake on the welfare of individually housed calves and found that those in all dietary regimes experienced elevated incidences of pneumonia and digestive illnesses. The lungs of all calves in the experiment showed signs of pneumonia at some point during the rearing period. As the calves were housed in individual stalls and denied solid food, the prevalence of pneumonia could have been linked to either of these factors.¹⁸³

Waltner-Toews *et al.* found that rearing calves outdoors in individual hutches, as opposed to indoors in adjacent pens, appeared to significantly decrease both diarrhea and pneumonia rates. There was no significant difference in the incidence of disease between individually housed and group-housed calves reared indoors.¹⁸⁴

Handling and Transport

Many calves, whether group-reared or tethered in a stall, endure the stress and discomfort of transport at least twice during their lifetimes. Most calves who will be raised for veal are trucked to livestock markets, where they are auctioned and then transported to the purchasing farm, though a small number may be sold directly to a veal facility.¹⁸⁵ After approximately 16-18 weeks¹⁸⁶ of fattening, the calves are transported to the slaughterhouse.

Studies have shown that transport can cause stress, fatigue, and injury. A study of the response of three-month-old calves to 18 hours of transport found reduced rumination and lying, excessive defecation and urination, and looseness of feces, all signs that the calves were stressed by both loading and vehicular movement.¹⁸⁷ Several studies have found that following transport calves show changes in appetite, signs of dehydration, and prolonged lying and sleeping behavior suggesting fatigue.^{188,189,190} Grigor *et al.* studied the effect of nine hours of road transport on young calves and reported high plasma cortisol responses, suggesting that one or more components of the transport were stressful.¹⁹¹ McCausland *et al.* examined 16,400 calves raised for veal and found that 50% had bruised stifle joints (akin to the human knee) after transport.¹⁹²

Although some of the transport-related welfare problems, such as dehydration and fatigue, may be exacerbated by long journeys, short trips are also stressful for calves due to loading and unloading. Grigor *et al.* reported higher heart rates both during and after a nine-hour journey, as well as greater plasma cortisol concentration and plasma creatine kinase activity immediately following the trip, compared with calves slaughtered on-farm who therefore were not transported.¹⁹³

Creatine kinase is a muscle-associated enzyme, and its concentration in blood can indicate the degree of physical exertion or muscle damage, such as bruising. The higher levels of this enzyme in the calves' bloodstream could indicate trauma at some point during the transport process—during loading, transport, and/or

unloading, and/or as a result of interaction among conspecifics. More bruised carcasses were observed in transported calves than those slaughtered on-farm.¹⁹⁴ These results echoed those of Van de Water *et al.*, who also found increased heart rate, plasma cortisol concentrations, and plasma creatine kinase activity in calves transported for 2-3 hours. The calves' behavior during loading—balking, reversing, and defecating—indicated the process was stressful. The researchers concluded that the entirety of the transport event, including loading, the journey, unloading, and lairage in a new environment, is stressful even if the journey portion is short, as plasma stress markers reached their maximal values within 30-60 minutes of the start of the journey.¹⁹⁵

Both severe and acute stresses are known to be immunosuppressant. As a consequence, increased disease incidence, particularly respiratory disease, has been observed in transported calves.¹⁹⁶ Staples and Hauge reportedly found that 60.3% of calves in their study transported before two weeks old fell ill during the following four weeks and 21.7% died. Pneumonia was identified as the greatest cause of death.¹⁹⁷ Similarly, Stephens reportedly recorded a mortality rate of 23% among calves transported for long distances during their first two weeks of life.¹⁹⁸ In a review of post-transport calf mortality, Knowles concluded that “young calves are not well adapted to cope with transport and marketing, often suffering relatively high rates of morbidity and mortality, both during, and in the few weeks immediately following transport” and recommended that calves not be taken to market until at least four weeks of age.¹⁹⁹

The effects of handling and transport are greater on individually housed calves than those group-reared. Trunkfield and Broom found that calves raised in stalls had higher blood cortisol concentrations after transport than calves of the same age who had been group-housed. The isolated calves had more difficulty walking and a significant increase in body temperature following transport. The researchers suggested that the increased temperature response was due in part to the trauma of walking for the first time in six months, particularly when boarding the ramp to the truck, as well as to the stress of mixing in close confinement with conspecifics for the first time.²⁰⁰ Lensink *et al.* attributed the significantly higher heart rate found among individually housed calves during transport to mixing with other calves, despite the fact that the individually housed animals could see and touch each other through the slats during rearing.²⁰¹

Slaughter

Given the heavy toll that transport takes on young calves, some arrive at the abattoir too weak to rise and walk, leaving them susceptible to abusive handling. In 2009, workers and the plant owner of the Bushway Packing plant in Grand Isle, Vermont were filmed shocking, kicking, and dragging downed calves. The plant owner also incorrectly used a captive bolt gun—meant for euthanizing calves not fit for slaughter—leaving them to die slowly.²⁰² In the press that followed the event, government officials stated that this kind of abuse was “inexcusable” and that abusive treatment is not representative of the industry as a whole,²⁰³ but this claim is unsubstantiated, and further investigations of cattle slaughter and auction facilities by The Humane Society of the United States suggest that abusive and neglectful treatment of cattle may not be uncommon.^{204,205}



Calves at Bushway Packing, Grand Isle, VT.
Photo by Humane Society of the United States.

According to the Humane Methods of Slaughter Act, U.S. federal law, calves should be “rendered insensible to pain by a single blow or gunshot or by electrical, chemical or other means that is rapid and effective, before being shackled, hoisted, thrown, cast or cut.”²⁰⁶

In a 2004 audit of five U.S. veal slaughter plants, slaughter expert Temple Grandin reported “some very bad problems,” including calves slipping on slick floors and excessive use of electric prods by truck drivers.²⁰⁷ According to Grandin, the most significant welfare problem was the shackling and hoisting of live calves for kosher slaughter at one plant. There, the vocalization of the calves was scored at 25%, compared to 3% or less at the plants that stunned calves—an increase Grandin attributes to the shackling and hoisting of fully sensible calves by one back leg.²⁰⁸

Stunning techniques in cattle, especially calves, can also present welfare problems.²⁰⁹ Large cattle are typically stunned using a captive bolt shot into the front of the head. When performed correctly, this method is effective,²¹⁰ causing a concussive blow to the head, tissue damage, and immediate²¹¹ or nearly immediate²¹² unconsciousness.

Electrical stunning is more commonly used on calves.²¹³ During electrical stunning, current is passed through the animal's brain to induce an epileptic state, as indicated by electroencephalogram recordings. In humans, there are no known instances of consciousness during an epileptic seizure of this type, and it is on this basis that animals are thought to be insensible during electrical stunning.²¹⁴ However, according to Anil *et al.*, electrical stunning of calves may produce shorter periods of unconsciousness and insensibility than with sheep or pigs.²¹⁵ Lambooy and Spanaard found that the shortest duration of unconsciousness following conventional (head-only) electrical stunning on calves was 21 seconds,²¹⁶ while others have reported periods of 45 seconds and 59 seconds.²¹⁷ As such, it can be assumed that calves may regain consciousness at some time between 22 and 60 seconds after head-only electrical stunning. Grandin surveyed several U.S. veal slaughterhouses and concluded:

A stunning method which produces either permanent or prolonged insensibility is essential for humane stunning of calves. I have observed calves reviving during bleeding in slaughter plants when conventional electric stunning was used. Calves may revive even if they are bled immediately after conventional electric stunning.²¹⁸

Grandin recommends the use of cardiac arrest stunning. In this method, electrodes are placed on either side of the calf's head and back or leg. Electric current runs through the brain and the heart simultaneously to produce unconsciousness and stop the heart, so the calf will not regain consciousness before being bled out. Sufficient amperage is important to ensure cardiac arrest. Blackmore and Petersen²¹⁹ reported that a head-to-leg stunner set at 2 amps for 5 seconds produced 100% cardiac arrest in calves, whereas 0.8 amps for 5 seconds was successful in just over half the calves. Grandin recommends a minimum amperage of 1.25, wetting the skin of the calves, and using good electrodes, but states that many variables will change amperage requirements and settings higher than 1.25 amps will be required in many slaughter plants.²²⁰ It is imperative that stunning equipment be routinely maintained and properly used,²²¹ or stunning may not be effective.

Conclusion

Presently, the customary veal production practices in the United States—namely housing in close, restrictive confinement, social isolation, deprivation of solid food, and provision of an iron-deficient diet—have been widely criticized on animal welfare grounds and are illegal throughout the European Union. The welfare of calves raised for veal can be dramatically improved with group housing on straw, provision of solid feed, and teat-feeding of milk, all of which have been shown to significantly reduce the suffering of calves during rearing. Closer scrutiny of calf slaughter is also in order, and abusive and neglectful treatment should not be tolerated. It is imperative that we continue to raise the bar for the care and treatment of veal calves, as legislators, food retailers, consumers, and the industry itself have already begun to demand.

¹ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 4, 2012.

² U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 4, 2012.

³ Keefe LM. 2008. The best of both worlds. *Meatingplace*, September, pp. 51-8.

⁴ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 4, 2012.

⁵ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 4, 2012.

-
- ⁶ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ⁷ Smith R. 2009. Veal producers ahead of schedule in housing transition. Feedstuffs, June 25. www.feedstuffs.com/ME2/dirmod.asp?sid=F4D1A9DFCD974EAD8CD5205E15C1CB42&nm=&type=news&mod=News&mid=A3D60400B4204079A76C4B1B129CB433&tier=3&nid=7D8D1E9F9C0B41B5A7DFED0F638DF9A0. Accessed March 8, 2012.
- ⁸ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ⁹ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 4, 2012.
- ¹⁰ The veal information gateway. 2011. Veal is the meat of a calf. www.veal.ca/home/. Accessed April 5, 2012.
- ¹¹ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ¹² Ohio Department of Agriculture. 2011. Ohio Livestock Care Standards, Bovine: Veal, Dairy, Beef. www.agri.ohio.gov/LivestockCareStandards/docs/OLCS%20Bovine%20-%20Final.pdf. Accessed April 5, 2012.
- Farmissues.com. 2012. Facts and figures about veal farming in Canada. Housing: where do veal calves live? www.farmissues.com/pages/factsVeal.php. Accessed April 4, 2012.
- ¹⁴ Ontario Veal Association. 2007. Where to buy Ontario veal (and other tips). www.ontariovealappeal.ca/buy.php. Accessed April 4, 2012.
- ¹⁵ House of Commons. 2004. Hansard. Written answers for 5 May 2004 (pt 4). www.publications.parliament.uk/pa/cm200304/cmhansrd/vo040505/text/40505w04.htm. Accessed April 5, 2012.
- ¹⁶ The Welfare of Farmed Animals (England) Regulations. 2000. Statutory Instrument 2000 No. 1870. www.opsi.gov.uk/SI/si2000/20001870.htm. Accessed April 5, 2012.
- ¹⁷ Council of Europe. 1997. Council Directive 97/2/EC of 20 January 1997 amending Directive 91/629/EEC laying down minimum standards for the protection of calves. http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-2-ec_en.pdf. Accessed April 5, 2012.
- ¹⁸ European Commission. 1997. Commission Decision of 24 February 1997 amending the Annex to Directive 91/629/EEC laying down minimum standards for the protection of calves (Text with EEA relevance) (97/182/EC). http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-182-ec_en.pdf. Accessed April 5, 2012.
- ¹⁹ European Commission. 1997. Commission Decision of 24 February 1997 amending the Annex to Directive 91/629/EEC laying down minimum standards for the protection of calves (Text with EEA relevance) (97/182/EC). http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-182-ec_en.pdf. Accessed April 5, 2012.
- ²⁰ Arizona Secretary of State's Office. 2006. Ballot Proposition Guide. Official Proposition 204 language. www.azsos.gov/election/2006/Info/PubPamphlet/Sun_Sounds/english/Prop204.htm. Accessed April 5, 2012.
- ²¹ Office of Gov. Bill Ritter, Jr. 2008. Gov. Ritter signs agriculture bills into law. Press release issued May 14. www.dehcc.org/2008Legislature/rittersigns051408.pdf. Accessed April 5, 2012.
- ²² State of Colorado. 2008. A Bill for an Act Concerning Requirements for Confinement of Specified livestock. www.statebillinfo.com/bills/bills/08/201_ren.pdf. Accessed April 18, 2012.
- ²³ California Health and Safety Code, Division 20, Chapter 13.8, Farm Animal Cruelty, Section 25990-25994. www.leginfo.ca.gov/cgi-bin/displaycode?section=hsc&group=25001-26000&file=25990-25994. Accessed April 5, 2012.
- ²⁴ California Secretary of State Debra Bowen. 2008. Statement of Vote, November 4, 2008, General Election. www.sos.ca.gov/elections/sov/2008_general/sov_complete.pdf. Accessed April 5, 2012.
- ²⁵ Hall C. 2008. Measure to provide better treatment of farm animals passes. Los Angeles Times, Nov. 5.
- ²⁶ Maine Public Law. 2009. Chapter 127, An act to prohibit cruel confinement of calves raised for veal and sows during gestation. www.mainelegislature.org/legis/bills/bills_124th/chapters/PUBLIC127.asp. Accessed April 5, 2012.

-
- ²⁷ Michigan Enrolled House Bill 5127 .2009. www.legislature.mi.gov/documents/2009-2010/billenrolled/House/pdf/2009-hNB-5127.pdf. Accessed April 5, 2012.
- ²⁸ Ohio Department of Agriculture. 2010. Statement from Ohio agriculture director regarding the passage of house bill 414 establishing the Ohio Livestock Care Standards Board. Press release issued March 24. http://ohiolivestockcarestandardsboard.gov/public_docs/news/3-24-10%20Directors%20Statement%20OLCSB.pdf. Accessed February 15, 2012.
- ²⁹ Ohio Department of Agriculture. 2011. Ohio Livestock Care Standards, Bovine: Veal, Dairy, Beef. www.agri.ohio.gov/LivestockCareStandards/docs/OLCS%20Bovine%20-%20Final.pdf. Accessed February 15, 2012.
- ³⁰ Marcelo P. 2012. New R.I. law bans cutting dairy-cow tails, raising pigs and calves in crates. Providence Journal, June 21. <http://news.providencejournal.com/breaking-news/2012/06/new-ri-law-bans.html>. Accessed June 27, 2012.
- ³¹ Smith R. 2007. Veal group housing approved. Feedstuffs, August 6, p. 3.
- ³² Keefe LM. 2012. Veal industry records progress in group housing. Meating Place, May 10. www.meatingplace.com/Industry/News/Details/32957. Accessed June 27, 2012.
- ³³ Salvage B. 2006. Revolutionizing the veal industry. Meat Processing, December, pp. 14-21.
- ³⁴ The Humane Society of the United States. 2007. Strauss Veal and Marcho Farms eliminating confinement by crate. www.humanesociety.org/assets/pdfs/farm/strauss_veal_marcho_farms.pdf. Accessed April 5, 2012.
- ³⁵ Salvage B. 2006. Revolutionizing the veal industry. Meat Processing, December, pp. 14-21.
- ³⁶ Strauss Group-Raised veal. www.straussbrands.com/groupraised/groupraised.html. Accessed March 12, 2012.
- ³⁷ Ketzenberger J. 2010. Hoosier farmers lead effort to raise veal humanely. The Indianapolis Star, August 16. www.indystar.com/article/20100817/LIVING07/8220303. Accessed March 12, 2012.
- ³⁸ Torres C. 2011. Producers Eager to Open Doors' at Farm Show. Lancaster Farming, December 31. www.lancasterfarming.com/-Producers-Eager-to--Open-Doors--at-Farm-Show-. Accessed March 12, 2012.
- ³⁹ Pew Commission on Industrial Farm Animal Production. 2008. Putting meat on the table: industrial farm animal production in America. www.ncifap.org/images/PCIFAPFin.pdf. Accessed April 5, 2012.
- ⁴⁰ Webster AJ, Saville C, Church BM, Gnanasakthy A, and Moss R. 1985. The effect of different rearing systems on the development of calf behaviour. British Veterinary Journal 141(3):249-64.
- ⁴¹ Andrighetto I, Gottardo F, Andreoli D, and Cozzi G. 1999. Effect of type of housing on veal calf growth performance, behaviour and meat quality. Livestock Production Science 57(2):137-45.
- ⁴² Sabbioni A, Berretti V, Bertocchi A, Zanon A, Soffiantini CS, and Superchi P. 2005. Effects of housing type on veal calf performance. Annali Facoltà Medicina Veterinaria di Parma XXV:111-30.
- ⁴³ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 5, 2012.
- ⁴⁴ Morisse JP, Cotte JP, Huonnic D, and Martrenchar A. 1999. Influence of dry feed supplements on different parameters of welfare in veal calves. Animal Welfare 8(1):43-52.
- ⁴⁵ Cozzi G, Gottardo F, Mattiello S, et al. 2002. The provision of solid feeds to veal calves: I. Growth performance, forestomach development, and carcass and meat quality. Journal of Animal Science 80(2):357-66.
- ⁴⁶ Van Putten G. 1982. Welfare in veal calf units. The Veterinary Record 111(19):437-40.
- ⁴⁷ U.S. Environmental Protection Agency. Ag 101. Lifestyle production phases. www.epa.gov/oecaagct/ag101/dairyphases.html. Accessed April 5, 2012.
- ⁴⁸ Flower FC and Weary DM. 2003. The effects of early separation on the dairy cow and calf. Animal Welfare 12(3):339-48.
- ⁴⁹ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ⁵⁰ American Veal Association. American veal industry fact sheet. <http://meatcuttersclub.activeboard.com/t45176320/american-veal-industry-fact-sheet/>. Accessed April 5, 2012.
- ⁵¹ Reinhardt V and Reinhardt A. 1981. Cohesive relationships in a cattle herd (Bos indicus). Behaviour 77:121-51.

-
- ⁵² Veissier I, Lamy D, and Le Neindre P. 1990. Social behaviour in domestic beef cattle when yearling calves are left with the cows for the next calving. *Applied Animal Behaviour Science* 27(3):193-200.
- ⁵³ Flower FC and Weary DM. 2003. The effects of early separation on the dairy cow and calf. *Animal Welfare* 12(3):339-48.
- ⁵⁴ Hudson SJ and Mullord MM. 1977. Investigations of maternal bonding in dairy cattle. *Applied Animal Ethology* 3(3):271-6.
- ⁵⁵ Lidfors LM. 1996. Behavioural effects of separating the dairy calf immediately or 4 days post-partum. *Applied Animal Behaviour Science* 49(3):269-83.
- ⁵⁶ Marchant-Forde JN, Marchant-Forde RM, and Weary DM. 2002. Responses of dairy cows and calves to each other's vocalisations after early separation. *Applied Animal Behaviour Science* 78(1):19-28.
- ⁵⁷ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 5, 2012.
- ⁵⁸ Flower FC and Weary DM. 2003. The effects of early separation on the dairy cow and calf. *Animal Welfare* 12(3):339-48.
- ⁵⁹ Flower FC and Weary DM. 2003. The effects of early separation on the dairy cow and calf. *Animal Welfare* 12(3):339-48.
- ⁶⁰ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ⁶¹ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13, citing: Stull CL and McMartin DA. 1992. Welfare parameters in veal calf production facilities. University of California, Davis, CA.
- ⁶² Stull CL and McDonough SP. 1994. Multidisciplinary approach to evaluating welfare of veal calves in commercial facilities. *Journal of Animal Science* 72(9):2518-24.
- ⁶³ Lidfors LM. 1996. Behavioural effects of separating the dairy calf immediately or 4 days post-partum. *Applied Animal Behaviour Science* 49(3):269-83.
- ⁶⁴ Weary DM and Chua B. 2000. Effects of early separation on the dairy cow and calf: 1. Separation at 6 h, 1 day, and 4 days after birth. *Applied Animal Behaviour Science* 69(3):177-88.
- ⁶⁵ Metz J and Metz JHM. 1986. Maternal influence on defecation and urination in the newborn calf. *Applied Animal Behaviour Science* 16(4):325-33.
- ⁶⁶ Van Metre DC, Tennant BC, and Whitlock RH. 2008. Infectious diseases of the gastrointestinal tract. In: Divers TJ and Peek SF (ed.s). 2008. *Rebhun's Diseases of Dairy Cattle*, 2nd Edition (St. Louis, MO: Saunders Elsevier, p. 203).
- ⁶⁷ Besser TE, Gay CC, and Pritchett L. 1991. Comparison of three methods of feeding colostrum to dairy calves. *Journal of the American Veterinary Medical Association* 198(3):419-22.
- ⁶⁸ Brignole TJ and Stott GH. 1980. Effect of suckling folloed by bottle feeding colostrum on immunoglobulin absorption and calf survival. *Journal of Dairy Science* 63:451-6.
- ⁶⁹ Godden S. 2008. Colostrum management for dairy calves. *Veterinary Clinics of North America* 24:19-39.
- ⁷⁰ Godden S. 2008. Colostrum management for dairy calves. *Veterinary Clinics of North America: Food Animal Practice* 24(1):19-39.
- ⁷¹ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ⁷² Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ⁷³ Hammell KL, Metz JHM, and Mekking P. 1988. Sucking behaviour of dairy calves fed milk ad libitum by bucket or teat. *Applied Animal Behaviour Science* 20(3):275-85.
- ⁷⁴ de Passillé AM. 2001. Sucking motivation and related problems in calves. *Applied Animal Behaviour Science* 72(3):175-87.
- ⁷⁵ de Passillé AM. 2001. Sucking motivation and related problems in calves. *Applied Animal Behaviour Science* 72(3):175-87.

- ⁷⁶ Jensen MB. 2003. The effects of feeding method, milk allowance and social factors on milk feeding behaviour and cross-sucking in group housed dairy calves. *Applied Animal Behaviour Science* 80(3):191-206.
- ⁷⁷ de Passillé AM. 2001. Sucking motivation and related problems in calves. *Applied Animal Behaviour Science* 72(3):175-87.
- ⁷⁸ Jensen MB. 2003. The effects of feeding method, milk allowance and social factors on milk feeding behaviour and cross-sucking in group housed dairy calves. *Applied Animal Behaviour Science* 80(3):191-206.
- ⁷⁹ Veissier I, de Passillé AM, Després G, et al. 2002. Does nutritive and non-nutritive sucking reduce other oral behaviors and stimulate rest in calves? *Journal of Animal Science* 80(10):2574-87.
- ⁸⁰ Veissier I, de Passillé AM, Després G, et al. 2002. Does nutritive and non-nutritive sucking reduce other oral behaviors and stimulate rest in calves? *Journal of Animal Science* 80(10):2574-87.
- ⁸¹ de Passillé AM. 2001. Sucking motivation and related problems in calves. *Applied Animal Behaviour Science* 72(3):175-87.
- ⁸² Haley DB, Rushen J, Duncan IJ, Widowski TM, and de Passillé AM. 1998. Effects of resistance to milk flow and the provision of hay on nonnutritive sucking by dairy calves. *Journal of Dairy Science* 81(8):2165-72.
- ⁸³ Gottardo F, Mattiello S, Cozzi G, et al. 2002. The provision of drinking water to veal calves for welfare purposes. *Journal of Animal Science* 80(9):2362-72.
- ⁸⁴ Council of Europe. 1997. Council Directive 97/2/EC of 20 January 1997 amending Directive 91/629/EEC laying down minimum standards for the protection of calves. http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-2-ec_en.pdf. Accessed April 5, 2012.
- ⁸⁵ European Commission. 1997. Commission Decision of 24 February 1997 amending the Annex to Directive 91/629/EEC laying down minimum standards for the protection of calves (Text with EEA relevance) (97/182/EC). http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-182-ec_en.pdf. Accessed April 5, 2012.
- ⁸⁶ Jensen MB. 2003. The effects of feeding method, milk allowance and social factors on milk feeding behaviour and cross-sucking in group housed dairy calves. *Applied Animal Behaviour Science* 80(3):191-206.
- ⁸⁷ Fraser AF and Broom DM. 1990. *Farm Animal Behaviour and Welfare*, Third Edition (London, U.K.: Bailliere Tindall).
- ⁸⁸ Veissier I, Ramirez de la Fe AR, and Pradel P. 1998. Nonnutritive oral activities and stress responses of veal calves in relation to feeding and housing conditions. *Applied Animal Behaviour Science* 57(1/2):35-49.
- ⁸⁹ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Fact_Sheets/Veal_from_Farm_to_Table/index.asp. Accessed April 5, 2012.
- ⁹⁰ Ngapo TM and Gariépy C. 2006. Factors affecting the meat quality of veal. *Journal of the Science of Food and Agriculture* 86(10):1412-31.
- ⁹¹ Ngapo TM and Gariépy C. 2006. Factors affecting the meat quality of veal. *Journal of the Science of Food and Agriculture* 86(10):1412-31.
- ⁹² Ngapo TM and Gariépy C. 2006. Factors affecting the meat quality of veal. *Journal of the Science of Food and Agriculture* 86(10):1412-31.
- ⁹³ Cozzi G, Gottardo F, Mattiello S, et al. 2002. The provision of solid feeds to veal calves: I. Growth performance, forestomach development, and carcass and meat quality. *Journal of Animal Science* 80(2):357-66.
- ⁹⁴ Fraser AF and Broom DM. 1990. *Farm Animal Behaviour and Welfare*, Third Edition (London, U.K.: Bailliere Tindall).
- ⁹⁵ Morisse JP, Cotte JP, Huonnic D, and Martrenchar A. 1999. Influence of dry feed supplements on different parameters of welfare in veal calves. *Animal Welfare* 8(1):43-52.
- ⁹⁶ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 6, 2012.
- ⁹⁷ Department for Environment, Food and Rural Affairs. 2003. Code of recommendations for the welfare of livestock: cattle (London, U.K.: Defra Publications). [www.agrowebcee.net/fileadmin/subnetwork/awsee/fawro/DOCS/Bovine/bovine1\(en\).pdf](http://www.agrowebcee.net/fileadmin/subnetwork/awsee/fawro/DOCS/Bovine/bovine1(en).pdf). Accessed April 6, 2012.
- ⁹⁸ European Commission. 1997. Commission Decision of 24 February 1997 amending the Annex to Directive 91/629/EEC laying down minimum standards for the protection of calves (Text with EEA relevance) (97/182/EC). http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-182-ec_en.pdf. Accessed April 6, 2012.

-
- ⁹⁹ Ngapo TM and Gariépy C. 2006. Factors affecting the meat quality of veal. *Journal of the Science of Food and Agriculture* 86(10):1412-31.
- ¹⁰⁰ Stull CL and McDonough SP. 1994. Multidisciplinary approach to evaluating welfare of veal calves in commercial facilities. *Journal of Animal Science* 72(9):2518-24.
- ¹⁰¹ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 6, 2012.
- ¹⁰² European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. *The EFSA Journal* 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹⁰³ European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to *The EFSA Journal* 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹⁰⁴ Welchman DD, Whelehan OP, and Webster AJ. 1988. Haematology of veal calves reared in different husbandry systems and the assessment of iron deficiency. *The Veterinary Record* 123(20):505-10.
- ¹⁰⁵ Reece WO and Hotchkiss DK. 1987. Blood studies and performance among calves reared by different methods. *Journal of Dairy Science* 70(8):1601-11.
- ¹⁰⁶ Morisse JP, Cotte JP, Huonnic D, and Martrenchar A. 1999. Influence of dry feed supplements on different parameters of welfare in veal calves. *Animal Welfare* 8(1):43-52.
- ¹⁰⁷ Cozzi G, Gottardo F, Mattiello S, et al. 2002. The provision of solid feeds to veal calves: I. Growth performance, forestomach development, and carcass and meat quality. *Journal of Animal Science* 80(2):357-66.
- ¹⁰⁸ Reece WO and Hotchkiss DK. 1987. Blood studies and performance among calves reared by different methods. *Journal of Dairy Science* 70(8):1601-11.
- ¹⁰⁹ Xiccato G, Trocino A, Queaque PI, Sartori A, and Carazzolo A. 2002. Rearing veal calves with respect to animal welfare: effects of group housing and solid feed supplementation on growth performance and meat quality. *Livestock Production Science* 75(3):269-80.
- ¹¹⁰ Reece WO and Hotchkiss DK. 1987. Blood studies and performance among calves reared by different methods. *Journal of Dairy Science* 70(8):1601-11.
- ¹¹¹ European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. *The EFSA Journal* 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹¹² Morisse JP, Cotte JP, Huonnic D, and Martrenchar A. 1999. Influence of dry feed supplements on different parameters of welfare in veal calves. *Animal Welfare* 8(1):43-52.
- ¹¹³ Veissier I, Ramirez de la Fe AR, and Pradel P. 1998. Nonnutritive oral activities and stress responses of veal calves in relation to feeding and housing conditions. *Applied Animal Behaviour Science* 57(1/2):35-49.
- ¹¹⁴ Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ¹¹⁵ European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to *The EFSA Journal* 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹¹⁶ Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ¹¹⁷ Morisse JP, Cotte JP, Huonnic D, and Martrenchar A. 1999. Influence of dry feed supplements on different parameters of welfare in veal calves. *Animal Welfare* 8(1):43-52.
- ¹¹⁸ Mattiello S, Canali E, Ferrante V, et al. 2002. The provision of solid feeds to veal calves: II. Behavior, physiology, and abomasal damage. *Journal of Animal Science* 80(2):367-75.
- ¹¹⁹ Veissier I, Ramirez de la Fe AR, and Pradel P. 1998. Nonnutritive oral activities and stress responses of veal calves in relation to feeding and housing conditions. *Applied Animal Behaviour Science* 57(1/2):35-49.

-
- ¹²⁰ Webster AJ, Saville C, Church BM, Gnanasakthy A, and Moss R. 1985. The effect of different rearing systems on the development of calf behaviour. *British Veterinary Journal* 141(3):249-64.
- ¹²¹ Wiepkema PR, van Hellemond KK, Roessingh P, and Romberg H. 1987. Behaviour and abomasal damage in individual veal calves. *Applied Animal Behaviour Science* 18(3/4):257-68.
- ¹²² Wiepkema PR, van Hellemond KK, Roessingh P, and Romberg H. 1987. Behaviour and abomasal damage in individual veal calves. *Applied Animal Behaviour Science* 18(3/4):257-68.
- ¹²³ Friend TH. 1991. Symposium: Response of animals to stress (Behavioral aspects of stress). *Journal of Dairy Science* 74(1):292-303.
- ¹²⁴ Wiepkema PR, van Hellemond KK, Roessingh P, and Romberg H. 1987. Behaviour and abomasal damage in individual veal calves. *Applied Animal Behaviour Science* 18(3/4):257-68.
- ¹²⁵ Wiepkema PR, van Hellemond KK, Roessingh P, and Romberg H. 1987. Behaviour and abomasal damage in individual veal calves. *Applied Animal Behaviour Science* 18(3/4):257-68.
- ¹²⁶ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 6, 2012.
- ¹²⁷ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. *Proceedings of the International Symposium in Le Mans, France, September 12-13*.
- ¹²⁸ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 6, 2012.
- ¹²⁹ Van Putten G. 1982. Welfare in veal calf units. *The Veterinary Record* 111(19):437-40.
- ¹³⁰ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹³¹ European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹³² European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36. www.efsa.europa.eu/en/efsajournal/doc/366.pdf. Accessed April 6, 2012.
- ¹³³ Stull CL and McDonough SP. 1994. Multidisciplinary approach to evaluating welfare of veal calves in commercial facilities. *Journal of Animal Science* 72(9):2518-24.
- ¹³⁴ Andrighetto I, Gottardo F, Andreoli D, and Cozzi G. 1999. Effect of type of housing on veal calf growth performance, behaviour and meat quality. *Livestock Production Science* 57(2):137-45.
- ¹³⁵ Smits AC and de Wilt JG. 1991. Group housing of veal calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 61-6).
- ¹³⁶ Webster AJ, Saville C, Church BM, Gnanasakthy A, and Moss R. 1985. The effect of different rearing systems on the development of calf behaviour. *British Veterinary Journal* 141(3):249-64.
- ¹³⁷ Wilson LL, Terosky TL, Stull CL, and Stricklin WR. 1999. Effects of individual housing design and size on behavior and stress indicators of special-fed Holstein veal calves. *Journal of Animal Science* 77(6):1341-7.
- ¹³⁸ Van Putten G. 1982. Welfare in veal calf units. *The Veterinary Record* 111(19):437-40.
- ¹³⁹ Le Neindre P. 1993. Evaluating housing systems for veal calves. *Journal of Animal Science* 71(5):1345-54.
- ¹⁴⁰ European Food Safety Authority. 2006. Scientific report on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36. www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_report_calveswelfare_en1.pdf. Accessed April 9, 2012.
- ¹⁴¹ European Commission. 1997. Commission Decision of 24 February 1997 amending the Annex to Directive 91/629/EEC laying down minimum standards for the protection of calves (Text with EEA relevance) (97/182/EC). http://ec.europa.eu/food/fs/aw/aw_legislation/calves/97-182-ec_en.pdf. Accessed April 9, 2012.

-
- ¹⁴² Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. Proceedings of the International Symposium in Le Mans, France, September 12-13.
- ¹⁴³ Department for Environment, Food and Rural Affairs. 2003. Code of recommendations for the welfare of livestock: cattle (London, U.K.: Defra Publications).
[www.agrowebcee.net/fileadmin/subnetwork/awsee/fawro/DOCS/Bovine/bovine1\(en\).pdf](http://www.agrowebcee.net/fileadmin/subnetwork/awsee/fawro/DOCS/Bovine/bovine1(en).pdf). Accessed April 9, 2012.
- ¹⁴⁴ European Food Safety Authority. 2006. Scientific report on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36.
www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_report_calveswelfare_en1.pdf. Accessed April 9, 2012.
- ¹⁴⁵ European Food Safety Authority. 2006. Scientific report on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36.
www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_report_calveswelfare_en1.pdf. Accessed April 9, 2012.
- ¹⁴⁶ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 9, 2012.
- ¹⁴⁷ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 9, 2012.
- ¹⁴⁸ Webster AJ, Saville C, Church BM, Gnanasakthy A, and Moss R. 1985. The effect of different rearing systems on the development of calf behaviour. *British Veterinary Journal* 141(3):249-64.
- ¹⁴⁹ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 9, 2012.
- ¹⁵⁰ Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ¹⁵¹ Reuters. 2000. Top New York restaurants stop serving white veal. July 6.
- ¹⁵² Jensen MB. 1999. Effects of confinement on rebounds of locomotor behaviour of calves and heifers, and the spatial preferences of calves. *Applied Animal Behaviour Science* 62(1):43-56.
- ¹⁵³ Jensen MB, Vestergaard KS, and Krohn CC. 1998. Play behaviour in dairy calves kept in pens: the effect of social contact and space allowance. *Applied Animal Behaviour Science* 56(2/4):97-108.
- ¹⁵⁴ Friend TH. 1991. Symposium: Response of animals to stress (Behavioral aspects of stress). *Journal of Dairy Science* 74(1):292-303.
- ¹⁵⁵ Dantzer R, Mormede P, Bluthe RM, and Soissons J. 1983. The effect of different housing conditions on behavioural and adrenocortical reactions in veal calves. *Reproduction Nutrition and Development* 23(3):501-8.
- ¹⁵⁶ Dellmeier GR, Friend TH, and Gbur EE. 1985. Comparison of four methods of calf confinement: II. Behavior. *Journal of Animal Science* 60(5):1102-9.
- ¹⁵⁷ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁵⁸ Flower FC and Weary DM. 2003. The effects of early separation on the dairy cow and calf. *Animal Welfare* 12(3):339-48.
- ¹⁵⁹ Reinhardt V and Reinhardt A. 1981. Cohesive relationships in a cattle herd (*Bos indicus*). *Behaviour* 77:121-51.
- ¹⁶⁰ European Food Safety Authority. 2006. Scientific report on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to The EFSA Journal 366:1-36.
www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_report_calveswelfare_en1.pdf. Accessed April 9, 2012.

-
- ¹⁶¹ Holm L, Jensen MB, and Jeppesen LL. 2002. Calves' motivation for access to two different types of social contact measured by operant conditioning. *Applied Animal Behaviour Science* 79(3):175-94.
- ¹⁶² Dellmeier GR, Friend TH, and Gbur EE. 1985. Comparison of four methods of calf confinement: II. Behavior. *Journal of Animal Science* 60(5):1102-9.
- ¹⁶³ European Food Safety Authority. 2006. Scientific opinion on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. *The EFSA Journal* 366:1-36.
www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_op_ej366_calveswelfare_en1.pdf. Accessed April 9, 2012.
- ¹⁶⁴ Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ¹⁶⁵ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁶⁶ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁶⁷ European Commission, Scientific Veterinary Committee, Animal Welfare Section. 1995. Report on the welfare of calves. Adopted November 9. http://ec.europa.eu/food/fs/sc/oldcomm4/out35_en.pdf. Accessed April 9, 2012.
- ¹⁶⁸ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62, citing: de Wilt JG. 1985. *Behaviour and Welfare of Veal Calves in Relation to Husbandry Systems*. Institute of Agricultural Engineering, Wageningen, The Netherlands (Thesis).
- ¹⁶⁹ Jensen MB, Vestergaard KS, Krohn CC, and Munksgaard L. 1997. Effect of single versus group housing and space allowance on responses of calves during open-field tests. *Applied Animal Behaviour Science* 54(2/3):109-21.
- ¹⁷⁰ European Food Safety Authority. 2006. Scientific report on the risks of poor welfare in intensive calf farming systems. An update of the Scientific Veterinary Committee report on the welfare of calves. Adopted May 24, 2006. Annex to *The EFSA Journal* 366:1-36.
www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/ahaw_report_calveswelfare_en1.pdf. Accessed April 9, 2012.
- ¹⁷¹ Broom DM. 1991. Needs and welfare of housed calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 23-31).
- ¹⁷² Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁷³ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁷⁴ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁷⁵ Dantzer R, Mormede P, Bluthé RM, and Soissons J. 1983. The effect of different housing conditions on behavioural and adrenocortical reactions in veal calves. *Reproduction Nutrition and Development* 23(3):501-8.
- ¹⁷⁶ Friend TH, Dellmeier GR, and Gbur EE. 1985. Comparison of four methods of calf confinement: I. Physiology. *Journal of Animal Science* 60(5):1095-101.
- ¹⁷⁷ Raussi S, Lensink BJ, Boissy A, Pyykkonen M, and Veissier I. 2003. The effect of contact with conspecifics and humans on calves' behaviour and stress responses. *Animal Welfare* 12(2):191-203.
- ¹⁷⁸ Stull C and Reynolds J. 2008. Calf welfare. *Veterinary Clinics of North America: Food Animal Practice* 24(1):191-203.
- ¹⁷⁹ Friend TH and Dellmeier GR. 1988. Common practices and problems related to artificially rearing calves: an ethological analysis. *Applied Animal Behaviour Science* 20(1/2):47-62.
- ¹⁸⁰ Webster AJF. 1991. Control of infectious disease in housed veal calves. In: Metz JHM and Groenestein CM (eds.), *New Trends in Veal Calf Production* (Wageningen, The Netherlands: EAAP Publications, pp. 103-11).
- ¹⁸¹ McDonough SP, Stull CL, and Osburn BI. 1994. Enteric pathogens in intensively reared veal calves. *American Journal of Veterinary Research* 55(11):1516-20.

-
- ¹⁸² McDonough SP, Stull CL, and Osburn BI. 1994. Enteric pathogens in intensively reared veal calves. *American Journal of Veterinary Research* 55(11):1516-20.
- ¹⁸³ McFarlane JM, Morris GL, Curtis SE, Simon J, and McGlone JJ. 1988. Some indicators of welfare of crated veal calves on three dietary iron regimens. *Journal of Animal Science* 66(2):317-25.
- ¹⁸⁴ Waltner-Toews D, Martin SW, and Meek AH. 1986. Dairy calf management, morbidity, and mortality in Ontario Holstein herds: III. Association of management with morbidity. *Preventive Veterinary Medicine* 4(2):137-58.
- ¹⁸⁵ Wilson LL, Stull CL, and Terosky TL. 1995. Veal perspectives to the year 2000: scientific advancements and legislation addressing veal calves in North America. *Proceedings of the International Symposium in Le Mans, France*, September 12-13.
- ¹⁸⁶ U.S. Department of Agriculture, Food Safety and Inspection Service. 2006. Veal from farm to table. www.fsis.usda.gov/Factsheets/Veal_from_Farm_to_Table/index.asp. Accessed April 9, 2012.
- ¹⁸⁷ Kent JE and Ewbank R. 1990. The behavioural response of 3-month-old calves to 18 hours road transportation. *Applied Animal Behaviour Science* 26(3):289.
- ¹⁸⁸ Kent JE and Ewbank R. 1990. The behavioural response of 3-month-old calves to 18 hours road transportation. *Applied Animal Behaviour Science* 26(3):289.
- ¹⁸⁹ Atkinson PJ. 1992. Investigation of the effects of transport and lairage on hydration state and resting behaviour of calves for export. *The Veterinary Record* 130(19):413-6.
- ¹⁹⁰ Grigor PN, Cockram MS, Steele WB, et al. 2001. Effects of space allowance during transport and duration of mid-journey lairage period on the physiological, behavioural and immunological responses of young calves during and after transport. *Animal Science* 73(pt. 2):341-60.
- ¹⁹¹ Grigor PN, Cockram MS, Steele WB, et al. 2001. Effects of space allowance during transport and duration of mid-journey lairage period on the physiological, behavioural and immunological responses of young calves during and after transport. *Animal Science* 73(pt. 2):341-60.
- ¹⁹² McCausland IP, Austin DF, and Dougherty R. 1977. Stifle bruising in bobby calves. *New Zealand Veterinary Journal* 25(3):71-2.
- ¹⁹³ Grigor PN, Cockram MS, Steele WB, et al. 2004. A comparison of the welfare and meat quality of veal calves slaughtered on the farm with those subjected to transportation and lairage. *Livestock Production Science* 91(3):219-28.
- ¹⁹⁴ Grigor PN, Cockram MS, Steele WB, et al. 2004. A comparison of the welfare and meat quality of veal calves slaughtered on the farm with those subjected to transportation and lairage. *Livestock Production Science* 91(3):219-28.
- ¹⁹⁵ Van de Water G, Verjans F, and Geers R. 2003. The effect of short distance transport under commercial conditions on the physiology of slaughter calves; pH and colour profiles of veal. *Livestock Production Science* 82(3):171-9.
- ¹⁹⁶ Trunkfield HR and Broom DM. 1990. The welfare of calves during handling and transport. *Applied Animal Behaviour Science* 28(1/2):135-52.
- ¹⁹⁷ Trunkfield HR and Broom DM. 1990. The welfare of calves during handling and transport. *Applied Animal Behaviour Science* 28(1/2):135-52, citing: Staples GE and Haugse CN. 1974. Losses in young calves after transportation. *British Veterinary Journal* 130:374-378.
- ¹⁹⁸ Knowles TG. 1995. A review of post transport mortality among younger calves. *The Veterinary Record* 137:406-7, citing: Stephens DB. 1982. In: Moss R (ed.), *Transport of Animals Intended for Breeding Production and Slaughter* (The Hague: Martinus Nijhoff, pp. 187-205).
- ¹⁹⁹ Knowles TG. 1995. A review of post transport mortality among younger calves. *The Veterinary Record* 137:406-7.
- ²⁰⁰ Trunkfield HR and Broom DM. 1990. The welfare of calves during handling and transport. *Applied Animal Behaviour Science* 28(1/2):135-52.
- ²⁰¹ Lensink BJ, Raussi S, Boivin X, Pyykkonen M, and Veissier I. 2001. Reactions of calves to handling depend on housing condition and previous experience with humans. *Applied Animal Behaviour Science* 70(3):187-99.
- ²⁰² Petition seeking closure of regulatory loophole facilitating mistreatment of calves too weak, sick or injured to stand and walk, *The Humane Society of the United States v. The Honorable Tom Vilsack and Jerold R. Mande*

(U.S. Department of Agriculture). www.fsis.usda.gov/PDF/Petition_HSUS_Humane_Handling.pdf. Accessed April 9, 2012.

²⁰³ Rampton, R. 2009. Vermont meat plant shut after USDA views footage. Reuters, October 30. www.reuters.com/article/2009/10/30/us-meat-humane-usda-idUSTRE59T4CO20091030. Accessed August 18, 2011.

²⁰⁴ Metzler NT. 2008. Humane Society releases new video of mistreated livestock. USA Today, May 7. www.usatoday.com/news/health/2008-05-07-3034539786_x.htm. Accessed March 8, 2012.

²⁰⁵ Gaouette N. 2008. Panel grills meatpacking chief on beef recall at Chino plant. Los Angeles Times, March 13. <http://articles.latimes.com/2008/mar/13/nation/na-hallmark13>. Accessed March 8, 2012.

²⁰⁶ Humane Methods of Slaughter Act. 7 United States Code. §§ 1901 et seq. <http://uscode.house.gov/download/pls/07C48.txt>. Accessed April 10, 2012.

²⁰⁷ Grandin T. 2004 Restaurant animal welfare audits of stunning and handling in federally inspected US and Canadian beef, veal, pork, lamb, and poultry slaughter plants. www.grandin.com/survey/2004.restaurant.audits.html. Accessed April 10, 2012.

²⁰⁸ Grandin T. 2004 Restaurant animal welfare audits of stunning and handling in federally inspected US and Canadian beef, veal, pork, lamb, and poultry slaughter plants. www.grandin.com/survey/2004.restaurant.audits.html. Accessed April 10, 2012.

²⁰⁹ Anil MH, McKinstry JL, Wotton SB, and Gregory NG. 1995. Welfare of calves: 1. Investigations into some aspects of calf slaughter. *Meat Science* 41(2):101-12.

²¹⁰ Anil MH, McKinstry JL, Wotton SB, and Gregory NG. 1995. Welfare of calves: 1. Investigations into some aspects of calf slaughter. *Meat Science* 41(2):101-12.

²¹¹ Svendsen O, Jensen SK, Karlsen LV, Svalastoga E, Jensen HE. 2008. Observations on newborn calves rendered unconscious with a captive bolt gun. *Veterinary Record* 162:90-2.

²¹² Gibson TJ, Johnson CB, Murrell JC, Mitchinson SL, Stafford KJ, and Mellor DJ. 2009. Electroencephalographic responses to concussive non-penetrative captive-bolt stunning in halothane-anaesthetised calves. *New Zealand Veterinary Journal* 57(2):90-5.

²¹³ Anil MH, McKinstry JL, Wotton SB, and Gregory NG. 1995. Welfare of calves: 1. Investigations into some aspects of calf slaughter. *Meat Science* 41(2):101-12.

²¹⁴ Gregory NG. 1998. *Animal Welfare and Meat Science* (Wallingford, UK: CABI Publishing, p.229).

²¹⁵ Anil MH, McKinstry JL, Wotton SB, and Gregory NG. 1995. Welfare of calves: 1. Investigations into some aspects of calf slaughter. *Meat Science* 41(2):101-12.

²¹⁶ Lambooy E and Spanjaard W. 1982. Electrical stunning of veal calves. *Meat Science* 6(1):15-25.

²¹⁷ Anil MH, McKinstry JL, Wotton SB, and Gregory NG. 1995. Welfare of calves: 1. Investigations into some aspects of calf slaughter. *Meat Science* 41(2):101-12.

²¹⁸ Grandin T. Cardiac arrest stunning of livestock and poultry with 1997 updates. www.grandin.com/humane/cardiac.arrest.html. Accessed April 10, 2012.

²¹⁹ Blackmore DK and Petersen GV. 1981. Stunning and slaughter of sheep and calves in New Zealand. *New Zealand Veterinary Journal* 29(6):99-102.

²²⁰ Grandin T. Cardiac arrest stunning of livestock and poultry with 1997 updates. www.grandin.com/humane/cardiac.arrest.html. Accessed April 10, 2012.

²²¹ Gregory NG. 1998. *Animal Welfare and Meat Science* (Wallingford, UK: CABI Publishing, pp.226, 230).

The Humane Society of the United States is the nation's largest animal protection organization—backed by 11 million Americans, or one of every 28. For more than a half-century, The HSUS has been fighting for the protection of all animals through advocacy, education, and hands-on programs. Celebrating animals and confronting cruelty. On the Web at humanesociety.org.