Gonçalves-de-Freitas, Eliane (2016) [Pain and fish welfare](#). *Animal Sentience* 3(22)
DOI: 10.51291/2377-7478.1058

Date of submission: 2015-11-02

Date of acceptance: 2015-12-13

This article has appeared in the journal *Animal Sentience*, a peer-reviewed journal on animal cognition and feeling. It has been made open access, free for all, by WellBeing International and deposited in the WBI Studies Repository. For more information, please contact wbisr-info@wellbeingintl.org.

Pain and fish welfare

Commentary on [Key](#) on Fish Pain

Eliane Gonçalves-de-Freitas

Departamento de Zoologia e Botânica
Universidade Estadual Paulista – UNESP

Abstract: The evolutionary approach of Key's (2016) target article, generically comparing humans with fish of all kinds, is simplistic. The author ignores published research on structural and molecular aspects of pain in fish. The target article reads more like a selective polemic against fish welfare than an even-handed analysis.

Keywords: nociception, opioids, welfare, evolution

Eliane Gonçalves de Freitas elianeg@ibilce.unesp.br,
Professor of Animal Behavior and Physiology at
University of the São Paulo State, does research on
fish social behavior, cognition, stress and welfare.
<http://lattes.cnpq.br/8774908691587814>

In the target article, Key (2016) draws attention to some interesting research challenges that remain open: how to measure animal consciousness and pain? Prominent omissions in Key's arguments, however, preclude accepting his conclusions. The evolutionary comparisons, for example, are presented in a simplistic way, based mostly on comparing fish (all kinds of fish) with just one mammal – humans. This is a phylogenetic mistake: As other commentators have pointed out, structures can evolve very differently, through convergent or divergent evolution.

A number of studies have shown the physiological framework for nociception and pain in fish (e.g., Sneddon, 2003a; Sneddon et al., 2003; Ashley et al., 2007), yet Key fails even to mention these important findings on the neuroarchitecture of fish pain. Other bioengineering components are likewise missing, especially molecular ones. Fish respond to opioids such as morphine (Sneddon, 2003b). This means fish have receptors for opioids, which subserve the defensive phase of pain in mammals (the absence of pain after injury, allowing animals to fight or flee). Evolutionarily, it is more plausible and parsimonious to infer that such an adaptive defensive mechanism was ancient and was preserved throughout vertebrate evolution.

One also cannot agree with Key's ominous suggestion that knowing that fish feel pain would be catastrophic. Catastrophic to whom? Not to the fish. A far more thoughtful (and empathic) approach is provided by Braithwaite's (2010) book "*Do fish feel pain?*" which is likewise absent from Key's lengthy list of references. In fact, it is hard not to get the impression that the purpose of Key's target article is less an objective attempt to assess the evidence for or against fish pain than a polemic against fish welfare. The dismissal of "benefit of the doubt" considerations as mere "anthropomorphism" is a case in point.

References

- Ashley, P.J., Sneddon, L.U. & McCrohan, C.R. (2007). Nociception in fish: stimulus-response properties of receptors on the head of trout *Oncorhynchus mykiss*. *Brain Research*, 1166, 47-54.
- Braithwaite, V. A. (2010). *Do Fish Feel Pain?* Oxford University Press, Oxford, United Kingdom.
- [Key, B. \(2016\)](#). Why fish do not feel pain. *Animal Sentience* 2016.003.
- Sneddon, L.U. (2003a). Trigeminal somatosensory innervation of the head of a teleost fish with particular reference to nociception. *Brain Research*, 972, 44–52.
- Sneddon, L.U. (2003b). The evidence for pain in fish: the use of morphine as an analgesic. *Applied Animal Behaviour Science*, 83, 153–162.
- Sneddon, L.U., Braithwaite, V.A. & Gentle, M.J. Do fishes have nociceptors? Evidence for the evolution of a vertebrate sensory system. *Proceedings of the Royal Society B: Biological Sciences*, 270, 1115-1121.